

GENERALIZED NUMERICAL RANGES AND QUANTUM ERROR CORRECTION

CHI-KWONG LI and YIU-TUNG POON

Communicated by Kenneth R. Davidson

ABSTRACT. For a noisy quantum channel, a quantum error correcting code of dimension k exists if and only if the joint rank- k numerical range associated with the error operators of the channel is non-empty. In this paper, geometric properties of the joint rank k -numerical range are obtained and their implications to quantum computing are discussed. It is shown that for a given k if the dimension of the underlying Hilbert space of the quantum states is sufficiently large, then the joint rank k -numerical range of operators is always star-shaped and contains the convex hull of the rank \widehat{k} -numerical range of the operators for sufficiently large \widehat{k} . In case the operators are infinite dimensional, the joint rank ∞ -numerical range of the operators is a convex set closely related to the joint essential numerical ranges of the operators.

KEYWORDS: *Quantum error correction, joint higher rank numerical range, joint essential numerical range, self-adjoint operator, Hilbert space.*

MSC (2000): 47A12, 15A60, 15A90, 81P68.

1. INTRODUCTION

In quantum computing, information is stored in *quantum bits*, abbreviated as *qubits*. Mathematically, a qubit is represented by a 2×2 rank one Hermitian matrix $Q = vv^*$, where $v \in \mathbb{C}^2$ is a unit vector. A state of N -qubits Q_1, \dots, Q_N is represented by their tensor products in M_n with $n = 2^N$. A *quantum channel* for states of N -qubits corresponds to a *trace preserving completely positive linear maps* $\Phi : M_n \rightarrow M_n$. By the structure theory of completely positive linear map [2], there are $T_1, \dots, T_r \in M_n$ with $\sum_{j=1}^r T_j^* T_j = I_n$ such that

$$(1.1) \quad \Phi(X) = \sum_{j=1}^r T_j X T_j^*.$$

In the context of quantum error correction, T_1, \dots, T_r are known as the *error operators*.

Let \mathbb{V} be a k -dimensional subspace of \mathbb{C}^n and P the orthogonal projection of \mathbb{C}^n onto \mathbb{V} . Then \mathbb{V} is a *quantum error correcting code* for the quantum channel Φ if there exists another trace preserving completely positive linear map $\Psi : M_n \rightarrow M_n$ such that $\Psi \circ \Phi(A) = A$ for all $A \in PM_nP$. By the results in [9], this happens if and only if there are scalars γ_{ij} with $1 \leq i, j \leq r$ such that

$$PT_i^*T_jP = \gamma_{ij}P.$$

Let \mathcal{P}_k be the set of rank k orthogonal projections in M_n . Define the *joint rank k -numerical range* of an m -tuple of matrices $\mathbb{A} = (A_1, \dots, A_m) \in M_n^m$ by

$$\Lambda_k(\mathbb{A}) = \{(a_1, \dots, a_m) \in \mathbb{C}^m : \text{there is } P \in \mathcal{P}_k \text{ such that } PA_jP = a_jP \text{ for } j = 1, \dots, m\}.$$

Then the quantum channel Φ defined in (1.1) has an error correcting code of k -dimension if and only if

$$\Lambda_k(T_1^*T_1, T_1^*T_2, \dots, T_r^*T_r) \neq \emptyset.$$

Evidently, $(a_1, \dots, a_m) \in \Lambda_k(\mathbb{A})$ if and only if there exists an $n \times k$ matrix U such that

$$U^*U = I_k, \quad \text{and} \quad U^*A_jU = a_jI_k \quad \text{for } j = 1, \dots, m.$$

Let $\mathbf{x}, \mathbf{y} \in \mathbb{C}^n$. Denote by $\langle \mathbb{A}\mathbf{x}, \mathbf{y} \rangle$ the vector $(\langle A_1\mathbf{x}, \mathbf{y} \rangle, \dots, \langle A_m\mathbf{x}, \mathbf{y} \rangle) \in \mathbb{C}^m$. Then $\mathbf{a} \in \Lambda_k(\mathbb{A})$ if and only if there exists an orthonormal set $\{\mathbf{x}_1, \dots, \mathbf{x}_k\}$ in \mathbb{C}^n such that $\langle \mathbb{A}\mathbf{x}_i, \mathbf{x}_i \rangle = \delta_{ij}\mathbf{a}$, where δ_{ij} is the Kronecker delta. When $k = 1$, $\Lambda_1(\mathbb{A})$ reduces to the (classical) *joint numerical range*

$$W(\mathbb{A}) = \{(\mathbf{x}^*A_1\mathbf{x}, \dots, \mathbf{x}^*A_m\mathbf{x}) : \mathbf{x} \in \mathbb{C}^n, \mathbf{x}^*\mathbf{x} = 1\}$$

of \mathbb{A} , which is quite well studied; see [11] and the references therein. It turns out that even for a single matrix $A \in M_n$, the study of $\Lambda_k(A)$ is highly non-trivial, and the results are useful in quantum computing, say, in constructing binary unitary channels; see [3], [4], [5], [6], [7], [14], [15], [18].

More generally, let $\mathcal{B}(\mathcal{H})$ be the algebra of bounded linear operators acting on a Hilbert space \mathcal{H} , which may be infinite dimensional. One can extend the definition of $\Lambda_k(A)$ to $A \in \mathcal{B}(\mathcal{H})$. If \mathcal{H} is infinite dimensional, one may allow $k = \infty$ by letting \mathcal{P}_k be the set of infinite rank orthogonal projections in $\mathcal{B}(\mathcal{H})$ in the definition; see [13], [16]. There are a number of reasons to consider rank k -numerical range of infinite dimensional operators. First, many quantum mechanical phenomena are better described using infinite dimensional Hilbert spaces. Also, a practical quantum computer must be able to handle a large number of qubits so that the underlying Hilbert space must have a very large dimension. We will also consider the joint rank k -numerical range of an m -tuple $\mathbb{A} = (A_1, \dots, A_m)$ of infinite dimensional operators A_1, \dots, A_m for positive integers k and $k = \infty$. It

is interesting to note that $\Lambda_\infty(\mathbb{A})$ has intimate connection with the *joint essential numerical range* of \mathbb{A} defined as

$$W_e(\mathbb{A}) = \bigcap \{ \text{cl}(W(\mathbb{A} + \mathbb{F})) : \mathbb{F} \in \mathcal{F}(\mathcal{H})^m \},$$

where $\mathcal{F}(\mathcal{H})$ denotes the set of finite rank operators in $\mathcal{B}(\mathcal{H})$ and $\text{cl}(\mathcal{S})$ denotes the closure of the set \mathcal{S} . Clearly, the joint essential numerical range is useful for the study of the joint behaviors of operators under perturbations of finite rank (or compact) operators.

The purpose of this paper is to study the joint rank k -numerical range of $\mathbb{A} = (A_1, \dots, A_m) \in \mathcal{B}(\mathcal{H})^m$. Understanding the properties of $\Lambda_k(\mathbb{A})$ is useful for constructing quantum error correcting codes and studying their properties such as their stability under perturbation.

Our paper is organized as follows. In Section 2, we present some basic properties of $\Lambda_k(\mathbb{A})$. Section 3 concerns the geometric properties of $\Lambda_k(\mathbb{A})$. We show that if $\dim \mathcal{H}$ is sufficiently large, then $\Lambda_k(\mathbb{A})$ is always star-shaped and contains the convex hull of $\Lambda_{\hat{k}}(\mathbb{A})$ for sufficiently large \hat{k} . In Section 4, we study the connection between $W_e(\mathbb{A})$, $\Lambda_k(\mathbb{A})$ and its closure $\text{cl}(\Lambda_k(\mathbb{A}))$. We show that $\Lambda_\infty(\mathbb{A})$ is always convex, and is a subset of the set of star centers of $\Lambda_k(\mathbb{A})$ for each positive integer k . We also show that

$$W_e(\mathbb{A}) = \bigcap_{k \geq 1} \text{cl}(\Lambda_k(\mathbb{A})).$$

Moreover, we obtain several equivalent formulations of $\Lambda_\infty(\mathbb{A})$, including

$$\Lambda_\infty(\mathbb{A}) = \bigcap \{ \Lambda_k(\mathbb{A} + \mathbb{F}) : \mathbb{F} \in \mathcal{F}(\mathcal{H})^m \}.$$

The results extend those in [1], [12].

Let $\mathcal{S}(\mathcal{H})$ be the real linear space of self-adjoint operators in $\mathcal{B}(\mathcal{H})$. Suppose

$$A_j = H_{2j-1} + iH_{2j} \quad \text{with } H_{2j-1}, H_{2j} \in \mathcal{S}(\mathcal{H}) \quad \text{for } j = 1, \dots, m.$$

Then $\Lambda_k(\mathbb{A}) \subseteq \mathbb{C}^m$ can be identified with $\Lambda_k(H_1, \dots, H_{2m}) \subseteq \mathbb{R}^{2m}$. Thus, we will focus on the joint rank k -numerical ranges of self-adjoint operators in our discussion.

2. BASIC PROPERTIES OF $\Lambda_k(\mathbb{A})$

PROPOSITION 2.1. *Suppose $\mathbb{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$, and $T = (t_{ij})$ is an $m \times n$ real matrix. If $B_j = \sum_{i=1}^m t_{ij} A_i$ for $j = 1, \dots, n$, then*

$$\{ \mathbf{a}T : \mathbf{a} \in \Lambda_k(\mathbb{A}) \} \subseteq \Lambda_k(\mathbb{B}).$$

Equality holds if $\{A_1, \dots, A_m\}$ is linearly independent and

$$\text{span} \{A_1, \dots, A_m\} = \text{span} \{B_1, \dots, B_n\}.$$

Proof. The set inclusion follows readily from definitions. Evidently, the equality holds if $n = m$ and T is invertible.

Suppose $\{A_1, \dots, A_m\}$ is linearly independent and

$$\text{span} \{A_1, \dots, A_m\} = \text{span} \{B_1, \dots, B_n\}.$$

First consider the special case when $A_i = B_i$ for $1 \leq i \leq m$. Then $T = [I_m | T_1]$ for some $m \times (n - m)$ matrix T_1 . Let $(b_1, \dots, b_n) \in \Lambda_k(\mathbb{B})$. Then there exists a rank k orthogonal projection P such that $PB_iP = b_iP$ for $i = 1, \dots, n$. Therefore, we have $(b_1, \dots, b_m) \in \Lambda_k(\mathbb{A})$ and for $1 \leq j \leq n$,

$$b_jP = PB_jP = P\left(\sum_{i=1}^m t_{ij}A_i\right)P = P\left(\sum_{i=1}^m t_{ij}B_i\right)P = \left(\sum_{i=1}^m t_{ij}b_i\right)P$$

implying that $b_j = \left(\sum_{i=1}^m t_{ij}b_i\right)$. Therefore, $(b_1, \dots, b_n) = (b_1, \dots, b_m)T$.

For the general case, by applying a permutation, if necessary, we may assume that $\{B_1, \dots, B_m\}$ is a basis of $\text{span} \{B_1, \dots, B_n\}$. Then there exists an $m \times m$ invertible matrix $S = (s_{ij})$ such that $A_j = \sum_{i=1}^m s_{ij}B_i$ for $j = 1, \dots, m$. For $1 \leq j \leq m$, we have

$$B_j = \sum_{i=1}^m t_{ij}A_i = \sum_{i=1}^m t_{ij}\left(\sum_{k=1}^m s_{ki}B_k\right) = \sum_{k=1}^m \left(\sum_{i=1}^m s_{ki}t_{ij}\right)B_k.$$

Therefore, $\sum_{i=1}^m s_{ki}t_{ij} = \delta_{kj}$ and $ST = [I_m | T_1]$ for some $m \times (n - m)$ matrix T_1 .

Hence, we have

$$\Lambda_k(\mathbb{B}) = \Lambda_k(B_1, \dots, B_m)[I | T_1] = \Lambda_k(B_1, \dots, B_m)ST = \Lambda_k(A_1, \dots, A_m)T. \quad \blacksquare$$

In view of the above proposition, in the study of the geometric properties of $\Lambda_k(\mathbb{A})$, we may always assume that A_1, \dots, A_m are linearly independent.

PROPOSITION 2.2. Let $\mathbb{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$, and let $k < \dim \mathcal{H}$.

(i) For any real vector $\boldsymbol{\mu} = (\mu_1, \dots, \mu_m)$,

$$\Lambda_k(A_1 - \mu_1 I, \dots, A_m - \mu_m I) = \Lambda_k(\mathbb{A}) - \boldsymbol{\mu}.$$

(ii) If $(a_1, \dots, a_m) \in \Lambda_k(\mathbb{A})$ then $(a_1, \dots, a_{m-1}) \in \Lambda_k(A_1, \dots, A_{m-1})$.

(iii) $\Lambda_{k+1}(\mathbb{A}) \subseteq \Lambda_k(\mathbb{A})$.

REMARK 2.3. By Proposition 2.2(i), we can replace A_j by $A_j - \mu_j I$ for $j = 1, \dots, m$, without affecting the geometric properties of $\Lambda_k(A_1, \dots, A_m)$.

Suppose $\dim \mathcal{H} = n < 2k - 1$ and $A_1 = \text{diag}(1, 2, \dots, n)$. Then $\Lambda_k(A_1) = \emptyset$. By Proposition 2.2(iii), we see that $\Lambda_k(\mathbb{A}) = \emptyset$ for any A_2, \dots, A_m . Thus, $\Lambda_k(\mathbb{A})$ can be empty if $\dim \mathcal{H}$ is small. However, a result of Knill, Laflamme and Viola [10] shows that $\Lambda_k(\mathbb{A})$ is non-empty if $\dim \mathcal{H}$ is sufficiently large. By modifying the proof of Theorem 3 in [10], we can get a slightly better bound in the following

proposition. The proof given here is essentially the same as that of Theorem 3 and 4 in [10], except for the choice of \mathbf{x}_1 . We include the details here for completeness.

PROPOSITION 2.4. *Let $\mathbb{A} \in \mathcal{S}(\mathcal{H})^m$. For $m \geq 1$ and $k > 1$. If*

$$\dim \mathcal{H} = n \geq (k - 1)(m + 1)^2,$$

then $\Lambda_k(\mathbb{A}) \neq \emptyset$.

Proof. We may assume that $\dim \mathcal{H} = n = (k - 1)(m + 1)^2$. Otherwise, replace each A_j by U^*A_jU for some U such that $U^*U = I_n$. Let

$$q = (m + 1)(k - 1) + 1.$$

Choose an eigenvector \mathbf{x}_1 of A_1 with $\|\mathbf{x}_1\| = 1$. Then choose a unit vector \mathbf{x}_2 orthogonal to $\mathbf{x}_1, A_2\mathbf{x}_1, \dots, A_m\mathbf{x}_1$. By the assumption on n , we can choose an orthonormal set $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_q\}$ of q vectors in \mathbb{C}^n such that for $1 < r \leq q$, \mathbf{x}_r is orthogonal to $A_j\mathbf{x}_i$ for all $1 \leq i < r$ and $1 \leq j \leq m$. Let X be the $n \times q$ matrix with \mathbf{x}_i as the i -th column. Then X^*A_jX is a diagonal matrix for $1 \leq j \leq m$. By Tverberg's Theorem [17], we can partition the set $\{i : 1 \leq i \leq q\}$ into k disjoint subset

$R_j, 1 \leq j \leq k$ such that $R = \bigcap_{j=1}^k \text{conv} \{ \langle \mathbb{A}\mathbf{x}_i, \mathbf{x}_i \rangle : i \in R_j \} \neq \emptyset$. Suppose $\mathbf{a} \in R$.

Then there exist non-negative numbers $t_{ij}, 1 \leq j \leq k, i \in R_j$ such that for all $1 \leq j \leq k, \sum_{i \in R_j} t_{ij} = 1$ and $\sum_{i \in R_j} t_{ij} \langle \mathbb{A}\mathbf{x}_i, \mathbf{x}_i \rangle = \mathbf{a}$. Let $\mathbf{y}_j = \sum_{i \in R_j} \sqrt{t_{ij}}\mathbf{x}_i$ for $1 \leq j \leq k$.

Then $\{\mathbf{y}_1, \dots, \mathbf{y}_k\}$ is orthonormal and $\langle \mathbb{A}\mathbf{y}_j, \mathbf{y}_j \rangle = \mathbf{a}$ for all $1 \leq j \leq k$. ■

PROPOSITION 2.5. *Suppose $\mathbb{A} \in \mathcal{S}(\mathcal{H})^m$ and $1 \leq r < k \leq \dim \mathcal{H}$. Let \mathcal{V}_r be the set of operators $X : \mathcal{H}_1^\perp \rightarrow \mathcal{H}$ such that $X^*X = I_{\mathcal{H}_1^\perp}$ for an r -dimensional subspace \mathcal{H}_1 of \mathcal{H} . Then*

$$(2.1) \quad \Lambda_k(\mathbb{A}) \subseteq \bigcap \{ \Lambda_{k-r}(X^*A_1X, \dots, X^*A_mX) : X \in \mathcal{V}_r \}$$

and

$$(2.2) \quad \text{conv} \Lambda_k(\mathbb{A}) \subseteq \text{conv} \left(\bigcap \{ \Lambda_{k-r}(X^*A_1X, \dots, X^*A_mX) : X \in \mathcal{V}_r \} \right).$$

Proof. Suppose $(a_1, \dots, a_m) \in \Lambda_k(\mathbb{A})$. Let \mathcal{H}_2 be a k -dimensional subspace and $V : \mathcal{H}_2 \rightarrow \mathcal{H}$ such that $V^*V = I_{\mathcal{H}_2}$ and $V^*A_jV = a_jI_k$ for $j = 1, \dots, m$. Let $X \in \mathcal{V}_r$ and $X^*X = I_{\mathcal{H}_1^\perp}$ for an r -dimensional subspace \mathcal{H}_1 of \mathcal{H} . Then

$$\mathcal{H}_0 = X^*(V(\mathcal{H}_2) \cap X(\mathcal{H}_1^\perp))$$

has dimension at least $s = k - r$. Let $U : \mathcal{H}_0 \hookrightarrow \mathcal{H}$ be given by $Ux = x$ for all $x \in \mathcal{H}_0$. Then we have $U^*U = I_{\mathcal{H}_0}$ and $U^*(X^*A_jX)U = a_jI_{\mathcal{H}_0}$ for $j = 1, \dots, m$. Thus, (2.1) holds, and the inclusion (2.2) follows. ■

Proposition 2.5 extends Proposition 4.8 of [13] corresponding to the case when $m = 2$. In such a case, the set inclusion (2.1) becomes a set equality if

$\dim \mathcal{H} < \infty$ or if (A_1, A_2) is a commuting pair, i.e., $A_1 + iA_2$ is normal; see Corollary 4.9 of [13]. The following example shows that the set equality in (2.1) may not hold even in the finite dimensional case if $m \geq 3$.

EXAMPLE 2.6. Let $B_1 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, $B_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, and $B_3 = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}$.

For $k > 1$, let $A_j = B_j \otimes I_k$ for $j = 1, 2, 3$.

(i) We have $\Lambda_k(A_1, A_2, A_3) = \Lambda_1(B_1, B_2, B_3) = \{\mathbf{a} \in \mathbb{R}^3 : \|\mathbf{a}\| = 1\}$, which is not convex.

(ii) If $r = k - 1$ and $X \in \mathcal{V}_r$ and $X^*X = I_{\mathcal{H}_1^\perp}$ for an r -dimensional subspace \mathcal{H}_1 , then $\dim \mathcal{H}_1^\perp = 2k - r = k + 1 \geq 3$ so that

$$\Lambda_{k-r}(X^*A_1X, X^*A_2X, X^*A_3X) = \Lambda_1(X^*A_1X, X^*A_2X, X^*A_3X)$$

is convex [11].

Consequently, $\bigcap\{\Lambda_{k-r}(X^*A_1X, X^*A_2X, X^*A_3X) : X \in \mathcal{V}_r\}$ is convex and cannot be equal to $\Lambda_k(A_1, A_2, A_3)$.

For $m > 3$, we can take A_1, A_2, A_3 as above and $A_j = 0_{2k}$ for $3 < j \leq m$. Then we have

$$\Lambda_k(A_1, \dots, A_m) \neq \bigcap\{\Lambda_{k-r}(X^*A_1X, \dots, X^*A_mX) : X \in \mathcal{V}_r\}.$$

To verify (i), suppose $U = \begin{pmatrix} U_1 \\ U_2 \end{pmatrix}$ is such that

$$U_1, U_2 \in M_k, \quad U^*U = U_1^*U_1 + U_2^*U_2 = I_k \quad \text{and} \quad U^*A_jU = a_jI_k.$$

Then $U_1^*U_1 - U_2^*U_2 = a_1I_k$. It follows that

$$U_1^*U_1 = (1 + a_1)I_k \quad \text{and} \quad U_2^*U_2 = (1 - a_1)I_k.$$

Thus, $U_1U_1^* = (1 + a_1)I_k$ and $U_2U_2^* = (1 - a_1)I_k$. As a result,

$$U_i^*U_jU_j^*U_i = (1 + a_1)(1 - a_1)I = U_i^*U_iU_j^*U_j \quad \text{for } (i, j) \in \{(1, 2), (2, 1)\},$$

and

$$\begin{aligned} (a_1^2 + a_2^2 + a_3^2)I_k &= \sum_{j=1}^3 (U^*A_jU)^2 \\ &= (U_1^*U_1 - U_2^*U_2)^2 + (U_1^*U_2 + U_2^*U_1)^2 + (iU_1^*U_2 - iU_2^*U_1)^2 \\ &= (U_1^*U_1 + U_2^*U_2)^2 = I_k. \end{aligned}$$

Thus, $\Lambda_k(A_1, A_2, A_3) \subseteq \{(a_1, a_2, a_3) \in \mathbb{R}^3 : a_1^2 + a_2^2 + a_3^2 = 1\}$.

Conversely, suppose $(a_1, a_2, a_3) \in \mathbb{R}^3$ such that $a_1^2 + a_2^2 + a_3^2 = 1$. Let

$$(\alpha, \beta) = \begin{cases} (0, 1) & \text{if } a_1 = -1, \\ \left(\frac{1+a_1, a_2-ia_3}{\sqrt{2(1+a_1)}}\right) & \text{otherwise.} \end{cases}$$

Let $U = \begin{pmatrix} \alpha I_k \\ \beta I_k \end{pmatrix}$. Direct computation shows that $U^*A_jU = a_jI_k$ for $1 \leq j \leq 3$.

By a similar argument or putting $k = 1$, we see that $\Lambda_1(B_1, B_2, B_3)$ has the same form.

It is natural to ask if the set equality in (2.2) can hold for $m > 2$. Also,

$$\begin{aligned} &\text{conv} \left(\bigcap \{ \Lambda_{k-r}(X^* A_1 X, \dots, X^* A_m X) : X \in \mathcal{V}_r \} \right) \\ &\subseteq \bigcap \{ \text{conv} \Lambda_{k-r}(X^* A_1 X, \dots, X^* A_m X) : X \in \mathcal{V}_r \}. \end{aligned}$$

It is interesting to determine whether the two sets are equal.

3. GEOMETRIC PROPERTIES OF $\Lambda_k(\mathbb{A})$

Let $\mathbb{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$. Much is known about $\Lambda_1(\mathbb{A})$; for example see [11] and its references. For instance, $\Lambda_1(\mathbb{A})$ is always convex if $m \leq 3$ unless $(\dim \mathcal{H}, m) = (2, 3)$; If $(\dim \mathcal{H}, m) = (2, 3)$ or $n > 1, m \geq 4$, there are examples of $\mathbb{A} \in \mathcal{S}(\mathcal{H})^m$ such that $\Lambda_1(\mathbb{A})$ is not convex. Furthermore, for any $A_1, A_2, A_3 \in \mathcal{S}(\mathcal{H})$ such that $\text{span} \{I, A_1, A_2, A_3\}$ has dimension 4, there is always an $A_4 \in \mathcal{S}(\mathcal{H})$ for which $\Lambda_1(A_1, \dots, A_4)$ is not convex.

In the following, we show that $\Lambda_k(\mathbb{A})$ is always star-shaped if $\dim \mathcal{H}$ is sufficiently large. Moreover, it always contains the convex hull of $\Lambda_{\widehat{k}}(\mathbb{A})$ for $\widehat{k} = (m+2)k$. If $k = 1$ and $m > 2$, we can lower the bound of the dimension of the Hilbert space to get the star-shapedness result. We begin with the following.

THEOREM 3.1. *Let $\mathbb{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$ and k be a positive integer. If $\Lambda_{\widehat{k}}(\mathbb{A}) \neq \emptyset$ for some $\widehat{k} \geq (m+2)k$, then $\Lambda_k(\mathbb{A})$ is star-shaped and contains the convex subset $\text{conv} \Lambda_{\widehat{k}}(\mathbb{A})$ so that every element in $\text{conv} \Lambda_{\widehat{k}}(\mathbb{A})$ is a star center of $\Lambda_k(\mathbb{A})$.*

Note that $\Lambda_{\widehat{k}}(\mathbb{A})$ may be empty if $\dim \mathcal{H}$ is small relative to \widehat{k} . Even if $\Lambda_{\widehat{k}}(\mathbb{A})$ is non-empty, it may be much smaller than its convex hull; for example, see Example 2.6. So, the conclusion in Theorem 3.1 is rather remarkable.

Proof. We may assume $\mathbf{a} = \mathbf{0} \in \Lambda_{\widehat{k}}(\mathbb{A})$. Then there exists Y such that $Y^*Y = I_{(m+2)k}$ and $Y^*A_jY = \mathbf{0}$ for all $1 \leq j \leq m$.

Let $\mathbf{b} \in \Lambda_k(\mathbb{A})$. Then there exists X such that $X^*X = I_k$ and

$$(X^* A_1 X, \dots, X^* A_m X) = (b_1 I_k, \dots, b_m I_k).$$

Suppose \mathcal{X} and \mathcal{Y} are the range spaces of X and Y , respectively. Then we have

$$\dim(\mathcal{Y} \cap (\mathcal{X} + A_1(\mathcal{X}) + \dots + A_m(\mathcal{X}))^\perp) \geq \dim \mathcal{Y} - \dim(\mathcal{X} + A_1(\mathcal{X}) + \dots + A_m(\mathcal{X})) \geq k.$$

Let \mathcal{Y}_1 be a k -dimensional subspace of $\mathcal{Y} \cap (\mathcal{X} + A_1(\mathcal{X}) + \dots + A_m(\mathcal{X}))^\perp$ and $\mathcal{Y}_2 = \mathcal{Y} \cap (\mathcal{X} + \mathcal{Y}_1)^\perp$. Set $Z = [X|Y_1|Y_2]$, where Y_i has columns forming an orthonormal basis of \mathcal{Y}_i for $i = 1, 2$. Then we have $Z^*Z = I_{(m+2)k}$ and for $1 \leq j \leq$

$m, Z^* A_j Z$ has the form

$$\begin{pmatrix} b_j I_k & \mathbf{0}_k & * \\ \mathbf{0}_k & \mathbf{0}_k & * \\ * & * & * \end{pmatrix}.$$

Let $C_j = b_j I_k \oplus \mathbf{0}_k$. For $t \in [0, 1]$, we have

$$t(b_1, \dots, b_m) \in \Lambda_k(C_1, \dots, C_m) \subseteq \Lambda_k(Z^* A_1 Z, \dots, Z^* A_m Z) \subseteq \Lambda_k(\mathbb{A}).$$

Clearly, $\Lambda_{\widehat{k}}(\mathbb{A}) \subseteq \Lambda_k(\mathbb{A})$. Since every element of $\Lambda_{\widehat{k}}(\mathbb{A})$ is a star center of $\Lambda_k(\mathbb{A})$ and the set of star centers of a star-shaped set is convex, we see that every element in $\text{conv } \Lambda_{\widehat{k}}(\mathbb{A})$ is a star center of $\Lambda_k(\mathbb{A})$. Hence, $\text{conv } \Lambda_{\widehat{k}}(\mathbb{A}) \subseteq \Lambda_k(\mathbb{A})$. ■

If $\dim \mathcal{H}$ is finite, then $\Lambda_k(\mathbb{A})$ is always closed. But this may not be the case if $\dim \mathcal{H}$ is infinite. Using Theorem 3.1, we can prove the star-shapedness of $\text{cl}(\Lambda_k(\mathbb{A}))$.

COROLLARY 3.2. *Let $\mathbb{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$ and k be a positive integer. If $\text{cl}(\Lambda_{\widehat{k}}(\mathbb{A})) \neq \emptyset$ for some $\widehat{k} \geq (m + 2)k$, then $\text{cl}(\Lambda_k(\mathbb{A}))$ is star-shaped and contains the convex subset $\text{conv cl}(\Lambda_{\widehat{k}}(\mathbb{A}))$ so that every element in $\text{conv cl}(\Lambda_{\widehat{k}}(\mathbb{A}))$ is a star center of $\text{cl}(\Lambda_k(\mathbb{A}))$.*

Proof. Suppose $\mathbf{a} \in \text{cl}(\Lambda_{\widehat{k}}(\mathbb{A}))$ and $\mathbf{b} \in \Lambda_k(\mathbb{A})$. Then for every ε there is $\tilde{\mathbf{a}} = (\tilde{a}_1, \dots, \tilde{a}_m) \in \Lambda_{\widehat{k}}(\mathbb{A})$ such that $\ell_1(\tilde{\mathbf{a}} - \mathbf{a}) < \varepsilon$. By Theorem 3.1, we see that the line segment joining $\tilde{\mathbf{a}}$ and \mathbf{b} lies in $\Lambda_k(\mathbb{A})$. Consequently, the line segment joining \mathbf{a} and \mathbf{b} lies in $\text{cl}(\Lambda_k(\mathbb{A}))$. The proof of the last assertion is similar to that of Theorem 3.1. ■

It is easy to see that a star center of $\Lambda_k(\mathbb{A})$ is also a star center of $\text{cl}(\Lambda_k(\mathbb{A}))$. However, the converse may not hold. The following example from Example 3.2 of [12] illustrates this.

EXAMPLE 3.3. Consider $\mathcal{H} = \ell^2$ with canonical basis $\{e_n : n \geq 1\}$. Let $\mathbb{A} = (A_1, \dots, A_4)$ with $A_1 = \text{diag}(1, 0, 1/3, 1/4, \dots)$, $A_2 = \text{diag}(1, 0) \oplus 0$,

$$A_3 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \oplus 0 \quad \text{and} \quad A_4 = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix} \oplus 0.$$

Then $(1, 1, 0, 0) \in W(\mathbb{A})$ and $(0, 0, 0, 0) \in W(\mathbb{A}) \cap W_e(\mathbb{A})$ is a star-center of the closure of $W(\mathbb{A})$. However, $(1/2, 1/2, 0, 0) \notin W(\mathbb{A})$ so that $(0, 0, 0, 0)$ is not a star-center of $W(\mathbb{A})$. In fact, $W(\mathbb{A})$ is not convex even though $\text{cl}(W(\mathbb{A}))$ is convex.

By Proposition 2.4, we see that $\Lambda_{\widehat{k}}(\mathbb{A})$ is non-empty if $\dim \mathcal{H}$ is sufficiently large. So, $\Lambda_k(\mathbb{A})$ is star-shaped and contains a convex set. The same comment also holds for $\text{cl}(\Lambda_k(\mathbb{A}))$. More specifically, we have the following.

THEOREM 3.4. *Let $\mathbb{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$. If*

$$\dim \mathcal{H} \geq ((m + 2)k - 1)(m + 1)^2,$$

then both $\Lambda_k(\mathbb{A})$ and $\text{cl}(\Lambda_k(\mathbb{A}))$ are star-shaped.

In Theorem 3.7, we will show that the classical joint numerical range is star-shaped with a much milder restriction on $\dim \mathcal{H}$ comparing with that in Theorem 3.4. To demonstrate this, we need two related results.

PROPOSITION 3.5. *Suppose $\dim \mathcal{H} = n$ and $\mathbb{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$ is such that $\{A_1, \dots, A_m\}$ is linearly independent. Assume that $\mathbf{0} \in \Lambda_{n-1}(\mathbb{A})$, i.e., there is a basis such that A_j has operator matrix $\begin{pmatrix} * & * \\ * & 0_{n-1} \end{pmatrix}$ for $j = 1, \dots, m$.*

(i) *If $m = 2n - 1$, then there is an invertible $S \in M_m(\mathbb{R})$ such that*

$$\Lambda_1(\mathbb{A}) = \left\{ (1 + u_1, u_2, \dots, u_m)S : u_1, \dots, u_m \in \mathbb{R}, \sum_{j=1}^m u_j^2 = 1 \right\}$$

so that $\Lambda_1(\mathbb{A})$ is not star-shaped.

(ii) *If $m < 2n - 1$, then $\Lambda_1(\mathbb{A})$ is star-shaped with $\mathbf{0}$ as a star center.*

Proof. (i) If $m = 2n - 1$, there is an invertible $m \times m$ real matrix $T = (t_{ij})$ such that for $B_j = \sum_{i=1}^m t_{ij}A_i$ for $j = 1, \dots, m$ with

$$\mathbb{B} = (B_1, \dots, B_m) = (E_{11}, E_{12} + E_{21}, -iE_{12} + iE_{21}, \dots, -iE_{1n} + iE_{n1}).$$

Let $\mathbf{x} = \mu(\cos t, \sin t(v_2 + iv_3), \dots, \sin t(v_{m-1} + iv_m))^t$ be a unit vector in \mathbb{C}^n such that $|\mu| = 1$, $t \in [0, \pi/2]$, and $v_2, \dots, v_m \in \mathbb{R}$ with $\sum_{j=2}^m v_j^2 = 1$. Then

$$\langle \mathbb{B}\mathbf{x}, \mathbf{x} \rangle = \left(\frac{1 + \cos(2t)}{2}, v_2 \sin(2t), v_2 \sin(2t), \dots, v_m \sin(2t) \right) = (1 + u_1, u_2, \dots, u_m)D,$$

where $D = [1/2] \oplus I_{n-1}$, $u_1 = \cos(2t)$ and $u_j = v_j \sin(2t)$ for $j = 2, \dots, m$. It follows that

$$\Lambda_1(\mathbb{B}) = \left\{ (1 + u_1, u_2, \dots, u_m)D : u_1, \dots, u_m \in \mathbb{R}, \sum_{j=1}^m u_j^2 = 1 \right\}.$$

By Proposition 2.1, $\Lambda_1(\mathbb{A}) = \{\mathbf{b}T^{-1} : \mathbf{b} \in \Lambda_1(\mathbb{B})\}$. The result follows.

(ii) Now, suppose $m < 2n - 1$. By adding more A_j , if necessary, we only need to consider the case when $m = 2n - 2$. Let \mathbf{v}_j be the row vector obtained by removing the first entry of the first row of A_j for $j = 1, \dots, m$.

Case 1. Suppose $\text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_m\}$ has real dimension $m - 1 = 2n - 3$. Then there is a unitary matrix of the form $U = [1] \oplus U_0 \in M_n$ such that the $(1, n)$ entry of U^*A_jU is real for $j = 1, \dots, m$. Hence, there is an invertible $m \times m$ real matrix $T = (t_{ij})$ such that for $B_j = \sum_{i=1}^m t_{ij}A_i$ for $j = 1, \dots, m$ with

$$\mathbb{B} = (B_1, \dots, B_m) = (E_{11}, E_{12} + E_{21}, -iE_{12} + iE_{21}, \dots, -iE_{1,n-1} + iE_{n-1,1}, E_{1n} + E_{n1}).$$

Suppose $\mathbf{b} \in \Lambda_1(\mathbb{B})$, $\mathbf{b} \neq \mathbf{0}$. Then there exists a unit vector $\mathbf{x} = \mu(u_0, u_1 + iu_2, \dots, u_{m-1} + iu_m)^t$ such that $|\mu| = 1$ and $u_0 > 0$, with

$$\mathbf{b} = \langle \mathbb{B}\mathbf{x}, \mathbf{x} \rangle = u_0(u_0, 2u_1, \dots, 2u_{m-1}).$$

For any $t \in (0, 1)$, we can choose a unit vector of the form $\mathbf{x}_t = \sqrt{t}(u_0, u_1 + iu_2, \dots, u_{m-1} + i\tilde{u}_m)^t$ with $t\tilde{u}_m^2 = 1 - \sum_{j=0}^{m-1} tu_j^2$ so that

$$\langle \mathbb{B}\mathbf{x}_t, \mathbf{x}_t \rangle = t\langle \mathbb{B}\mathbf{x}, \mathbf{x} \rangle.$$

Case 2. Suppose $\{\mathbf{v}_1, \dots, \mathbf{v}_m\}$ has real dimension $m = 2n - 2$. Then there is an invertible $m \times m$ real matrix $T = (t_{ij})$ such that

$$\begin{aligned} \mathbb{B} &= (B_1, \dots, B_m) = \left(\sum_{i=1}^m t_{i1}A_i, \dots, \sum_{i=1}^m t_{im}A_i \right) \\ &= (a_1E_{11}, \dots, a_mE_{11}) + (E_{12} + E_{21}, iE_{12} + iE_{21}, \dots, E_{1n} + E_{n1}, -iE_{1n} + iE_{n1}) \end{aligned}$$

with $a_1, \dots, a_m \in \mathbb{R}$. Suppose $\mathbf{b} \in \Lambda_1(\mathbb{B})$, $\mathbf{b} \neq \mathbf{0}$. Then there exists a unit vector $\mathbf{x} = \mu(u_0, u_1 + iu_2, \dots, u_{m-1} + iu_m)^t$ such that $|\mu| = 1$ and $u_0 > 0$, with

$$\mathbf{b} = \langle \mathbb{B}\mathbf{x}, \mathbf{x} \rangle = u_0(a_1u_0 + 2u_1, a_2u_0 + 2u_2, \dots, a_mu_0 + 2u_m).$$

For any $t \in (0, 1)$, consider a vector of the form

$$\mathbf{x}_\xi = (\xi u_0, w_1 + iw_2, w_3 + iw_4, \dots, w_{m-1} + iw_m)^t,$$

where $\xi \geq t$ and $w_j = a_j u_0(t - \xi^2)/(2\xi) + tu_j/\xi$ for $j = 1, \dots, m$. Then

$$\xi u_0(a_j \xi u_0 + 2w_j) = tu_0(a_j u_0 + 2u_j), \quad j = 1, \dots, m,$$

so that

$$\langle \mathbb{B}\mathbf{x}_\xi, \mathbf{x}_\xi \rangle = t\langle \mathbb{B}\mathbf{x}, \mathbf{x} \rangle.$$

If $\xi = \sqrt{t}$, then $\mathbf{x}_\xi = \sqrt{t}(u_0, u_1 + iu_2, \dots, u_{m-1} + iu_m)^t$ has norm less than 1; if $\xi \rightarrow \infty$, then $\|\mathbf{x}_\xi\| \geq |\xi u_0| \rightarrow \infty$. Thus, there is $\xi > t$ such that \mathbf{x}_ξ is a unit vector satisfying $\langle \mathbb{B}\mathbf{x}_\xi, \mathbf{x}_\xi \rangle = t\langle \mathbb{B}\mathbf{x}, \mathbf{x} \rangle$. So, $\Lambda_1(\mathbb{B})$ is star-shaped with $\mathbf{0}$ as a star center. By Proposition 2.1, $\Lambda_1(\mathbb{A}) = \{\mathbf{b}T^{-1} : \mathbf{b} \in \Lambda_1(\mathbb{B})\}$. The result follows. ■

THEOREM 3.6. *Let $\mathbb{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$. If $\Lambda_{\widehat{k}}(\mathbb{A}) \neq \emptyset$ for some $\widehat{k} > (m + 1)/2$, then $\Lambda_1(\mathbb{A})$ is star-shaped and contains $\text{conv } \Lambda_{\widehat{k}}(\mathbb{A})$ such that every element in $\text{conv } \Lambda_{\widehat{k}}(\mathbb{A})$ is a star center of $\Lambda_1(\mathbb{A})$.*

Proof. We may assume $\mathbf{a} = \mathbf{0} \in \Lambda_{\widehat{k}}(\mathbb{A})$ with $\widehat{k} > (m + 1)/2$. Suppose $\mathbf{x} \in \mathcal{H}$ is a unit vector and $\mathbf{b} = \langle \mathbb{A}\mathbf{x}, \mathbf{x} \rangle \in \Lambda_1(\mathbb{A})$. Suppose X is such that $X^*X = I_{\widehat{k}}$ and $X^*A_jX = 0_{\widehat{k}}$ for $j = 1, \dots, m$. Let Y be such that $Y^*Y = I_{\widehat{k}+1}$, and the range space of Y contains the range space of X and \mathbf{x} . Suppose $\mathbb{B} = (B_1, \dots, B_m) = (Y^*A_1Y, \dots, Y^*A_mY)$. Then we may assume that B_j has the form $\begin{pmatrix} * & * \\ * & 0_{\widehat{k}} \end{pmatrix}$ for $j = 1, \dots, m$. Clearly, $\text{span}\{B_1, \dots, B_m\}$ has dimension at most $m < 2\widehat{k} - 1$. By Proposition 3.5, the line segment joining $\mathbf{0}$ and \mathbf{b} lies entirely in $\Lambda_1(\mathbb{B}) \subseteq \Lambda_1(\mathbb{A})$. Thus, $\mathbf{0}$ is a star center of $\Lambda_1(\mathbb{A})$. Since the set of star centers of $\Lambda_1(\mathbb{A})$ is convex, we see that every element in $\text{conv } \Lambda_{\widehat{k}}(\mathbb{A})$ is a star center of $\Lambda_1(\mathbb{A})$. ■

THEOREM 3.7. Let $\mathbb{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$. If

$$\dim \mathcal{H} \geq \left\lceil \frac{m+1}{2} \right\rceil (m+1)^2,$$

then $\Lambda_1(\mathbb{A})$ is star-shaped.

Proof. Let $\widehat{k} = \lceil (m+1)/2 \rceil + 1 > (m+1)/2$. Then $\dim \mathcal{H} \geq (\widehat{k}-1)(m+1)^2$ and $\Lambda_{\widehat{k}}(\mathbb{A}) \neq \emptyset$ by Proposition 2.4. The result then follows from Theorem 3.6. ■

4. RESULTS ON $\Lambda_\infty(\mathbb{A})$

In this section, we always assume that \mathcal{H} has infinite dimension. Denote by \mathcal{P}_∞ the set of infinite rank orthogonal projections in $\mathcal{S}(\mathcal{H})$. For $\mathbb{A} \in \mathcal{S}(\mathcal{H})^m$ let

$$\Lambda_\infty(\mathbb{A}) = \{(\gamma_1, \dots, \gamma_m) \in \mathbb{R}^m : \text{there is } P \in \mathcal{P}_\infty \text{ such that } PA_iP = \gamma_iP \text{ for all } 1 \leq i \leq m\}.$$

By the result in Section 3, we have the following.

PROPOSITION 4.1. Suppose $\mathbb{A} \in \mathcal{S}(\mathcal{H})^m$, where \mathcal{H} is infinite-dimensional. Then $\Lambda_k(\mathbb{A})$ is star-shaped for each positive integer k . Moreover, if $\mathbf{a} \in \Lambda_\infty(\mathbb{A})$, then \mathbf{a} is a star center for $\Lambda_k(\mathbb{A})$ for every positive integer k .

When $m = 2$, it was conjectured in [16] and confirmed in [13] that

$$\Lambda_\infty(A_1, A_2) = \bigcap_{k \geq 1} \Lambda_k(A_1, A_2);$$

in Theorem 4 of [1], it was proven that

$$\Lambda_\infty(A_1, A_2) = \bigcap \{W(A_1 + F_1, A_2 + F_2) : F_1, F_2 \in \mathcal{S}(\mathcal{H}) \cap \mathcal{F}(\mathcal{H})\}.$$

In the following, we extend the above results to $\Lambda_\infty(A_1, \dots, A_m)$ for $m > 2$. Moreover, we show that $\Lambda_\infty(\mathbb{A}) = \bigcap_{k \geq 1} S_k(\mathbb{A})$, where $S_k(\mathbb{A})$ is the set of star centers of $\Lambda_k(\mathbb{A})$. Hence, $\Lambda_\infty(\mathbb{A})$ is always convex.

THEOREM 4.2. Suppose $\mathbb{A} \in \mathcal{S}(\mathcal{H})^m$, where \mathcal{H} is infinite-dimensional. For each $k \geq 1$, let $S_k(\mathbb{A})$ be the set of star-center of $\Lambda_k(\mathbb{A})$. Then

$$(4.1) \quad \Lambda_\infty(\mathbb{A}) = \bigcap_k S_k(\mathbb{A}) = \bigcap_k \Lambda_k(\mathbb{A}) = \bigcap \{W(\mathbb{A} + \mathbb{F}) : \mathbb{F} \in \mathcal{S}(\mathcal{H})^m \cap \mathcal{F}(\mathcal{H})^m\}.$$

Consequently, $\Lambda_\infty(\mathbb{A})$ is convex.

Proof. It follows from definitions and Theorem 3.1 that

$$\Lambda_\infty(\mathbb{A}) \subseteq \bigcap_k S_k(\mathbb{A}) \subseteq \bigcap_k \Lambda_k(\mathbb{A}).$$

We are going to prove that $\bigcap_k \Lambda_k(\mathbb{A}) \subseteq \bigcap \{W(\mathbb{A} + \mathbb{F}) : \mathbb{F} \in \mathcal{S}(\mathcal{H})^m \cap \mathcal{F}(\mathcal{H})^m\}$.

Suppose $\mathbb{F} = (F_1, \dots, F_m) \in \mathcal{S}(\mathcal{H})^m \cap \mathcal{F}(\mathcal{H})^m$ and $K = \sum_{i=1}^m \text{rank}(F_i) + 1$. Let $\boldsymbol{\mu} = (\mu_1, \dots, \mu_m) \in \bigcap_k \Lambda_k(\mathbb{A})$. Then there exists a rank K orthogonal projection P such that $PA_jP = \mu_j P$ for $1 \leq j \leq m$. Let

$$\begin{aligned} \mathcal{H}_0 &= \text{range } P \cap \ker F_1 \cap \ker F_2 \cap \dots \cap \ker F_m \\ &= \text{range } P \cap (\text{range } F_1 + \text{range } F_2 + \dots + \text{range } F_m)^\perp. \end{aligned}$$

Then $\dim \mathcal{H}_0 \geq 1$. Let \mathbf{x} be a unit vector in \mathcal{H}_0 . Then we have $\langle (\mathbb{A} + \mathbb{F})\mathbf{x}, \mathbf{x} \rangle = \langle \mathbb{A}\mathbf{x}, \mathbf{x} \rangle = \boldsymbol{\mu}$. Therefore, $\boldsymbol{\mu} \in W(\mathbb{A} + \mathbb{F})$. Hence, we have $\bigcap_k \Lambda_k(\mathbb{A}) \subseteq \bigcap \{W(\mathbb{A} + \mathbb{F}) : \mathbb{F} \in \mathcal{S}(\mathcal{H})^m \cap \mathcal{F}(\mathcal{H})^m\}$.

Next, we prove that $\bigcap \{W(\mathbb{A} + \mathbb{F}) : \mathbb{F} \in \mathcal{S}(\mathcal{H})^m \cap \mathcal{F}(\mathcal{H})^m\} \subseteq \Lambda_\infty(\mathbb{A})$. Suppose

$$\boldsymbol{\mu} \in \bigcap \{W(\mathbb{A} + \mathbb{F}) : \mathbb{F} \in \mathcal{S}(\mathcal{H})^m \cap \mathcal{F}(\mathcal{H})^m\}.$$

By Remark 2.3, we may assume that $\boldsymbol{\mu} = \mathbf{0}$. Then $\mathbf{0} \in W(\mathbb{A})$ and there exists a unit vector \mathbf{x}_1 such that $\langle \mathbb{A}\mathbf{x}_1, \mathbf{x}_1 \rangle = \mathbf{0}$. Suppose we have chosen an orthonormal set of vectors $\{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ such that $\langle \mathbb{A}\mathbf{x}_i, \mathbf{x}_j \rangle = \mathbf{0}$ for all $1 \leq i, j \leq n$. Let \mathcal{H}_0 be the subspace spanned by

$$\{\mathbf{x}_i : 1 \leq i \leq n\} \cup \{A_j\mathbf{x}_i : 1 \leq i \leq n, 1 \leq j \leq m\}$$

and P the orthogonal projection of \mathcal{H} onto \mathcal{H}_0 . Suppose

$$\mathbb{B} = ((I - P)A_1(I - P)|_{\mathcal{H}_0^\perp}, \dots, (I - P)A_m(I - P)|_{\mathcal{H}_0^\perp}).$$

Let $\mathbf{b} = (b_1, \dots, b_m)$ be a star-center of $W(\mathbb{B})$. Then

$$\mathbf{b}I_{\mathcal{H}_0} \oplus \mathbb{B} = (b_1P + (I - P)A_1(I - P), \dots, b_mP + (I - P)A_m(I - P)) = \mathbb{A} + \mathbb{F}$$

for some $\mathbb{F} \in \mathcal{S}(\mathcal{H})^m \cap \mathcal{F}(\mathcal{H})^m$. Therefore, $\mathbf{0} \in W(\mathbf{b}I_{\mathcal{H}_0} \oplus \mathbb{B})$. Hence, there exists a unit vector $\mathbf{x} \in \mathcal{H}$ such that $\mathbf{0} = \langle (\mathbb{A} + \mathbb{F})\mathbf{x}, \mathbf{x} \rangle$. Let $\mathbf{x} = \mathbf{y} + \mathbf{z}$, where $\mathbf{y} \in \mathcal{H}_0$ and $\mathbf{z} \in \mathcal{H}_0^\perp$. Then $\|\mathbf{y}\|^2 + \|\mathbf{z}\|^2 = \|\mathbf{x}\|^2 = 1$. If $\mathbf{z} = \mathbf{0}$, then $\mathbf{0} = \mathbf{b} \in W(\mathbb{B})$. If $\mathbf{z} \neq \mathbf{0}$, then by Proposition 4.1, we have

$$\mathbf{0} = \langle (\mathbb{A} + \mathbb{F})\mathbf{x}, \mathbf{x} \rangle = \|\mathbf{y}\|^2 \mathbf{b} + \|\mathbf{z}\|^2 \left\langle \mathbb{B} \left(\frac{\mathbf{z}}{\|\mathbf{z}\|} \right), \left(\frac{\mathbf{z}}{\|\mathbf{z}\|} \right) \right\rangle \in W(\mathbb{B}).$$

So there exists a unit vector $\mathbf{x}_{n+1} \in \mathcal{H}_0^\perp$ such that

$$\mathbf{0} = \langle (\mathbb{A} + \mathbb{F})\mathbf{x}_{n+1}, \mathbf{x}_{n+1} \rangle = \langle \mathbb{B}\mathbf{x}_{n+1}, \mathbf{x}_{n+1} \rangle = \langle \mathbb{A}\mathbf{x}_{n+1}, \mathbf{x}_{n+1} \rangle.$$

Hence, inductively, we can choose an orthonormal sequence of vectors $\{\mathbf{x}_n\}_{n=1}^\infty$ such that

$$\langle \mathbb{A}\mathbf{x}_i, \mathbf{x}_j \rangle = \mathbf{0} \quad \text{for all } i, j.$$

Thus, we have

$$\Lambda_\infty(\mathbb{A}) \subseteq \bigcap_k S_k(\mathbb{A}) \subseteq \bigcap_k \Lambda_k(\mathbb{A}) \subseteq \bigcap \{W(\mathbb{A} + \mathbb{F}) : \mathbb{F} \in \mathcal{S}(\mathcal{H})^m \cap \mathcal{F}(\mathcal{H})^m\} \subseteq \Lambda_\infty(\mathbb{A}).$$

Since $S_k(\mathbb{A})$ is convex for all $k \geq 1$, the last statement follows. ■

The last equality in (4.1) establishes a relationship between $\Lambda_\infty(\mathbb{A})$ and the joint numerical ranges of finite rank perturbations of \mathbb{A} . The following result gives an extension.

THEOREM 4.3. *Suppose $\mathbb{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$. Let $n \geq 1$ and $\mathbb{F}_0 \in \mathcal{S}(\mathcal{H})^m \cap \mathcal{F}(\mathcal{H})^m$. Then the following sets are equal:*

- (i) $\Lambda_\infty(\mathbb{A})$.
- (ii) $\Lambda_\infty(\mathbb{A} + \mathbb{F}_0)$.
- (iii) $\bigcap \{ \Lambda_n(\mathbb{A} + \mathbb{F}) : \mathbb{F} \in \mathcal{F}(\mathcal{H})^m \cap \mathcal{S}(\mathcal{H})^m \}$.
- (iv) $\bigcap_{k \geq 1} \left(\bigcap \{ \Lambda_k(\mathbb{A} + \mathbb{F}) : \mathbb{F} \in \mathcal{F}(\mathcal{H})^m \cap \mathcal{S}(\mathcal{H})^m \} \right)$.

Proof. By Theorem 4.2, we have

$$\begin{aligned} \Lambda_\infty(\mathbb{A} + \mathbb{F}_0) &= \bigcap \{ W(\mathbb{A} + \mathbb{F}_0 + \mathbb{F}) : \mathbb{F} \in \mathcal{S}(\mathcal{H})^m \cap \mathcal{F}(\mathcal{H})^m \} \\ &= \bigcap \{ W(\mathbb{A} + \mathbb{F}) : \mathbb{F} \in \mathcal{S}(\mathcal{H})^m \cap \mathcal{F}(\mathcal{H})^m \} = \Lambda_\infty(\mathbb{A}). \end{aligned}$$

This proves the equality of the sets in (i) and (ii). For the equality of the sets of (i) and (iii), let $\mathbb{F} \in \mathcal{F}(\mathcal{H})^m \cap \mathcal{S}(\mathcal{H})^m$. Then we have

$$\Lambda_\infty(\mathbb{A}) = \Lambda_\infty(\mathbb{A} + \mathbb{F}) \subseteq \Lambda_n(\mathbb{A} + \mathbb{F}) \subseteq W(\mathbb{A} + \mathbb{F}).$$

It follows that

$$\begin{aligned} \Lambda_\infty(\mathbb{A}) &\subseteq \bigcap \{ \Lambda_n(\mathbb{A} + \mathbb{F}) : \mathbb{F} \in \mathcal{S}(\mathcal{H})^m \cap \mathcal{F}(\mathcal{H})^m \} \\ &\subseteq \bigcap \{ W(\mathbb{A} + \mathbb{F}) : \mathbb{F} \in \mathcal{S}(\mathcal{H})^m \cap \mathcal{F}(\mathcal{H})^m \} = \Lambda_\infty(\mathbb{A}). \end{aligned}$$

The equivalence of (i) and (iv) follows immediately. ■

Recall that \mathcal{V}_r is the set of $X : \mathcal{H}_1^\perp \rightarrow \mathcal{H}$ such that $\dim \mathcal{H}_1 = r$ and $X^*X = I_{\mathcal{H}_1^\perp}$. Then $X^*\mathbb{A}X$ is a compression of \mathbb{A} to \mathcal{H}_1^\perp . The next result is an analog to Theorem 4.3 for $\Lambda_k(X^*\mathbb{A}X)$.

THEOREM 4.4. *Suppose $\mathbb{A} = (A_1, \dots, A_m) \in \mathcal{S}(\mathcal{H})^m$. Let $n, r_0 \geq 1$ and $X_0 \in \mathcal{V}_{r_0}$. Then for $X^*\mathbb{A}X = (X^*A_1X, \dots, X^*A_mX)$, the following sets are equal:*

- (i) $\Lambda_\infty(\mathbb{A})$.
- (ii) $\Lambda_\infty(X_0^*\mathbb{A}X_0)$.
- (iii) $\bigcap \left\{ \Lambda_n(X^*\mathbb{A}X) : X \in \bigcup_{r \geq 1} \mathcal{V}_r \right\}$.
- (iv) $\bigcap_{k \geq 1} \left(\bigcap \left\{ \Lambda_k(X^*\mathbb{A}X) : X \in \bigcup_{r \geq 1} \mathcal{V}_r \right\} \right)$.

Proof. By (4.1) and (2.1), we have

$$\Lambda_\infty(\mathbb{A}) = \bigcap_k \Lambda_k(\mathbb{A}) = \bigcap_k \Lambda_{k+r}(\mathbb{A}) \subseteq \bigcap_k \Lambda_k(X_0^*\mathbb{A}X_0) = \Lambda_\infty(X_0^*\mathbb{A}X_0) \subseteq \Lambda_\infty(\mathbb{A}).$$

This proves the equality of the sets in (i) and (ii). For the equality of the sets of (i) and (iii), we will first show that

$$(4.2) \quad \bigcap \left\{ \Lambda_1(X^* \mathbb{A} X) : X \in \bigcup_{r \geq 1} \mathcal{V}_r \right\} \subseteq \Lambda_\infty(\mathbb{A}).$$

Let $\mu \in \bigcap \left\{ \Lambda_1(X^* \mathbb{A} X) : X \in \bigcup_{r \geq 1} \mathcal{V}_r \right\}$. By Remark 2.3, we may assume that $\mu = \mathbf{0}$.

Then there exists a unit vector \mathbf{x}_1 such that $\langle \mathbb{A} \mathbf{x}_1, \mathbf{x}_1 \rangle = \mathbf{0}$. Suppose we have chosen an orthonormal set of vectors $\{\mathbf{x}_1, \dots, \mathbf{x}_N\}$ such that $\langle \mathbb{A} \mathbf{x}_i, \mathbf{x}_j \rangle = \mathbf{0}$ for all $1 \leq i, j \leq N$. Let \mathcal{H}_1 be the subspace spanned by

$$\{\mathbf{x}_i : 1 \leq i \leq N\} \cup \{A_j \mathbf{x}_i : 1 \leq i \leq N, 1 \leq j \leq m\}$$

and $X : \mathcal{H}_1^\perp \rightarrow \mathcal{H}$ be given by $X(\mathbf{v}) = \mathbf{v}$ for all $\mathbf{v} \in \mathcal{H}_1^\perp$. Then $\mathbf{0} \in \Lambda_1(X^* \mathbb{A} X)$. So there exists a unit vector $\mathbf{x}_{N+1} \in \mathcal{H}_1^\perp$ such that

$$\mathbf{0} = \langle (X^* \mathbb{A} X) \mathbf{x}_{N+1}, \mathbf{x}_{N+1} \rangle = \langle \mathbb{A} \mathbf{x}_{N+1}, \mathbf{x}_{N+1} \rangle.$$

Inductively, we can find an orthonormal sequence $\{\mathbf{x}_i\}$ in \mathcal{H} such that $\langle \mathbb{A} \mathbf{x}_i, \mathbf{x}_j \rangle = \mathbf{0}$ for all i, j . Hence, $\mathbf{0} \in \Lambda_\infty(\mathbb{A})$.

To continue the proof of the equality of the sets of (i) and (iii). Let $X \in \bigcup_{r \geq 1} \mathcal{V}_r$.

Then we have

$$\Lambda_\infty(\mathbb{A}) = \Lambda_\infty(X^* \mathbb{A} X) \subseteq \Lambda_n(X^* \mathbb{A} X) \subseteq \Lambda_1(X^* \mathbb{A} X).$$

It follows that

$$\Lambda_\infty(\mathbb{A}) \subseteq \bigcap \left\{ \Lambda_\infty(X^* \mathbb{A} X) : X \in \bigcup_{r \geq 1} \mathcal{V}_r \right\} \subseteq \bigcap \left\{ \Lambda_1(X^* \mathbb{A} X) : X \in \bigcup_{r \geq 1} \mathcal{V}_r \right\} \subseteq \Lambda_\infty(\mathbb{A}).$$

The equality of the sets of (i) and (iv) follows immediately. ■

Recall that the joint essential numerical range of $\mathbb{A} \in \mathcal{S}(\mathcal{H})^m$ is defined by

$$W_e(\mathbb{A}) = \bigcap \{ \text{cl}(W(\mathbb{A} + \mathbb{F})) : \mathbb{F} \in \mathcal{S}(\mathcal{H})^m \cap \mathcal{F}(\mathcal{H})^m \}.$$

Using the last two theorems, we have the following.

COROLLARY 4.5. *Let $\mathbb{A} \in \mathcal{S}(\mathcal{H})^m$, where \mathcal{H} is infinite dimensional. Denote by $\tilde{\mathcal{S}}_k(\mathbb{A})$ the set of star centers of $\text{cl}(\Lambda_k(\mathbb{A}))$. Then*

$$W_e(\mathbb{A}) = \bigcap_{k \geq 1} \text{cl}(\Lambda_k(\mathbb{A})) = \bigcap_{k \geq 1} \tilde{\mathcal{S}}_k(\mathbb{A}).$$

In addition, let $n \geq 1$ and $\mathbb{F}_0 \in \mathcal{S}(\mathcal{H})^m \cap \mathcal{F}(\mathcal{H})^m$. Moreover, let \mathbb{V} be a finite dimensional subspace of \mathcal{H} and $X_0 : \mathbb{V}^\perp \rightarrow \mathcal{H}$ such that $X_0^ X_0 = I_{\mathbb{V}^\perp}$. Then the following sets are equal:*

- (i) $W_e(\mathbb{A})$.
- (ii) $W_e(\mathbb{A} + \mathbb{F}_0)$.
- (iii) $W_e(X_0^* \mathbb{A} X_0)$.
- (iv) $\bigcap \{ \text{cl}(\Lambda_n(\mathbb{A} + \mathbb{F})) : \mathbb{F} \in \mathcal{F}(\mathcal{H})^m \cap \mathcal{S}(\mathcal{H})^m \}$.

- (v) $\bigcap \left\{ \text{cl}(\Lambda_n(X^* \mathbb{A} X)) : X \in \bigcup_{r \geq 1} \mathcal{V}_r \right\}$.
- (vi) $\bigcap_{k \geq 1} (\bigcap \{ \text{cl}(\Lambda_k(\mathbb{A} + \mathbb{F})) : \mathbb{F} \in \mathcal{F}(\mathcal{H})^m \cap \mathcal{S}(\mathcal{H})^m \})$.
- (vii) $\bigcap_{k \geq 1} \left(\bigcap \left\{ \text{cl}((\Lambda_k(X^* \mathbb{A} X)) : X \in \bigcup_{r \geq 1} \mathcal{V}_r \right\} \right)$.

Let $A \in \mathcal{S}(\mathcal{H})$ and k be a positive integer. Denote by \mathcal{U}_k the set of $X : \mathbb{C}^k \rightarrow \mathcal{H}$ such that $X^* X = I_k$ and

$$\lambda_k(A) = \sup \{ \min \sigma(X^* A X) : X \in \mathcal{U}_k \},$$

where $\sigma(B)$ is the spectrum of the operator B . For $\mathbf{c} = (c_1, \dots, c_m) \in \mathbb{R}^m$ and $\mathbb{A} \in \mathcal{S}(\mathcal{H})^m$, let $\mathbf{c} \cdot \mathbb{A} = \sum_{i=1}^m c_i A_i$. Define

$$\Omega_\infty(\mathbb{A}) = \bigcap_{\mathbf{c} \in \mathbb{R}^m} \{ \mathbf{a} \in \mathbb{R}^m : \mathbf{c} \cdot \mathbf{a} \leq \lambda_k(\mathbf{c} \cdot \mathbb{A}) \text{ for all } k \geq 1 \}.$$

The following extends Theorem 2.1 of [13].

THEOREM 4.6. *Let $\mathbb{A} \in \mathcal{S}(\mathcal{H})^m$, where \mathcal{H} is infinite dimensional. Then*

$$\Lambda_\infty(\mathbb{A}) \subseteq \Omega_\infty(\mathbb{A}) = W_e(\mathbb{A}).$$

Proof. For $k \geq 1$, let

$$\Omega_k(\mathbb{A}) = \bigcap_{\mathbf{c} \in \mathbb{R}^m} \{ \mathbf{a} \in \mathbb{R}^m : \mathbf{c} \cdot \mathbf{a} \leq \lambda_k(\mathbf{c} \cdot \mathbb{A}) \}.$$

Clearly, $\Omega_\infty(\mathbb{A}) = \bigcap_{k \geq 1} \Omega_k(\mathbb{A})$. Suppose $k \geq 1$ and $\mathbf{a} = (a_1, \dots, a_m) \in \Lambda_k(\mathbb{A})$. Then there exists $P \in \mathcal{P}_k$ such that $PA_jP = a_jP$ for all $1 \leq j \leq m$. For every $\mathbf{c} \in \mathbb{R}^m$, we have $P(\mathbf{c} \cdot \mathbb{A})P = (\mathbf{c} \cdot \mathbf{a})P$. Hence,

$$\mathbf{c} \cdot \mathbf{a} = \lambda_k(\mathbf{c} \cdot PAP) = \lambda_k(P(\mathbf{c} \cdot \mathbb{A})P) \leq \lambda_k(\mathbf{c} \cdot \mathbb{A}).$$

Therefore, $\Lambda_k(\mathbb{A}) \subseteq \Omega_k(\mathbb{A})$. Since $\Omega_k(\mathbb{A})$ is closed, we have $\text{cl}(\Lambda_k(\mathbb{A})) \subseteq \Omega_k(\mathbb{A})$. Hence,

$$\Lambda_\infty(\mathbb{A}) = \bigcap_{k \geq 1} \Lambda_k(\mathbb{A}) \subseteq \bigcap_{k \geq 1} \text{cl}(\Lambda_k(\mathbb{A})) = W_e(\mathbb{A}) \subseteq \bigcap_{k \geq 1} \Omega_k(\mathbb{A}) = \Omega_\infty(\mathbb{A}).$$

To show that $\Omega_\infty(\mathbb{A}) \subseteq W_e(\mathbb{A})$. Suppose $\mathbf{a} \in \Omega_k(\mathbb{A})$. By Remark 2.3, we may assume that $\mathbf{a} = \mathbf{0}$. So, $\lambda_k(\mathbf{c} \cdot \mathbb{A}) \geq 0$ for all $k \geq 1$ and $\mathbf{c} \in \mathbb{R}^m$. For $\mathbb{F} = (F_1, \dots, F_m) \in \mathcal{S}(\mathcal{H})^m \cap \mathcal{F}(\mathcal{H})^m$, let $K = \sum_{i=1}^m \text{rank}(F_i) + 1$. Then

$$\lambda_1(\mathbf{c} \cdot (\mathbb{A} + \mathbb{F})) \geq \lambda_{K+1}(\mathbf{c} \cdot \mathbb{A}) \geq 0 \text{ and } \lambda_1(-(\mathbf{c} \cdot (\mathbb{A} + \mathbb{F}))) \geq \lambda_{K+1}(-\mathbf{c} \cdot \mathbb{A}) \geq 0.$$

Therefore, $\mathbf{c} \cdot \mathbf{0} = 0 \in \text{cl}(W(\mathbf{c} \cdot (\mathbb{A} + \mathbb{F}))) = \mathbf{c} \cdot \text{cl}(W(\mathbb{A} + \mathbb{F}))$. Hence, $\mathbf{c} \cdot \mathbf{0} \in \mathbf{c} \cdot W_e(\mathbb{A})$ for all $\mathbf{c} \in \mathbb{R}^m$. By the convexity of $W_e(\mathbb{A})$, we have $\mathbf{0} \in W_e(\mathbb{A})$. ■

EXAMPLE 4.7. For $n \geq 1$, let $B_n = \begin{bmatrix} \frac{1}{n} & 0 \\ 0 & -\frac{1}{n} \end{bmatrix}$, $C_n = \begin{bmatrix} \frac{1}{n} & 0 \\ 0 & 0 \end{bmatrix}$, $A_1 = \bigoplus_{n=1}^{\infty} B_n$, $A_2 = \bigoplus_{n=1}^{\infty} C_n$ and $\mathbb{A} = (A_1, A_2)$. Then $(0, 0) \in \Omega_{\infty}(\mathbb{A})$ but $\Lambda_{\infty}(\mathbb{A}) = \emptyset$.

Acknowledgements. Li is an honorary professor of the University of Hong Kong and an honorary professor of the Taiyuan University of Technology. His research was partially supported by an NSF grant and the William and Mary Pulmeri Award. Poon's research was partially supported by an NSF grant.

REFERENCES

- [1] J. ANDERSON, J.G. STAMPFLI, Compressions and commutators, *Israel J. Math.* **10**(1971), 433–441.
- [2] M.D. CHOI, Completely positive linear maps on complex matrices, *Linear Algebra Appl.* **10**(1975), 285–290.
- [3] M.D. CHOI, M. GIESINGER, J.A. HOLBROOK, D.W. KRIBS, Geometry of higher-rank numerical ranges, *Linear and Multilinear Algebra* **56**(2008), 53–64.
- [4] M.D. CHOI, J.A. HOLBROOK, D.W. KRIBS, K. ŻYCZKOWSKI, Higher-rank numerical ranges of unitary and normal matrices, *Operators and Matrices* **1**(2007), 409–426.
- [5] M.D. CHOI, D.W. KRIBS, K. ŻYCZKOWSKI, Higher-rank numerical ranges and compression problems, *Linear Algebra Appl.* **418**(2006), 828–839.
- [6] M.D. CHOI, D.W. KRIBS, K. ŻYCZKOWSKI, Quantum error correcting codes from the compression formalism, *Rep. Math. Phys.* **58**(2006), 77–91.
- [7] H.L. GAU, C.K. LI, P.Y. WU, Higher-rank numerical ranges and dilations, *J. Operator Theory*, to appear.
- [8] K.E. GUSTAFSON, D.K.M. RAO, *Numerical Ranges: The Field of Values of Linear Operators and Matrices*, Springer, New York 1997.
- [9] E. KNILL, R. LAFLAMME, Theory of quantum error correcting codes, *Phys. Rev. A* **55**(1997), 900–911.
- [10] E. KNILL, R. LAFLAMME, L. VIOLA, Theory of quantum error correction for general noise, *Phys. Rev. Lett.* **84**(2000), 2525–2528.
- [11] C.K. LI, Y.T. POON, Convexity of the joint numerical range, *SIAM J. Matrix Anal. Appl.* **21**(1999), 668–678.
- [12] C.K. LI, Y.T. POON, The joint essential numerical range of operators: Convexity and related results, *Studia Math.* **194**(2009), 91–104.
- [13] C.K. LI, Y.T. POON, N.S. SZE, Higher rank numerical ranges and low rank perturbations of quantum channels, *J. Math. Anal. Appl.* **348**(2008), 843–855.
- [14] C.K. LI, Y.T. POON, N.S. SZE, Condition for the higher rank numerical range to be non-empty, *Linear and Multilinear Algebra* **57**(2009), 365–368.

- [15] C.K. LI, N.K. SZE, Canonical forms, higher rank numerical ranges, totally isotropic subspaces, and matrix equations, *Proc. Amer. Math. Soc.* **136**(2008), 3013–3023.
- [16] R.A. MARTINEZ-AVENDANO, Higher-rank numerical range in infinite-dimensional Hilbert space, *Operators and Matrices* **2**(2008), 249–264.
- [17] H. TVERBERG, A generalization of Radon's theorem, *J. London Math. Soc.* **41**(1966), 123–128.
- [18] H. WOERDEMAN, The higher rank numerical range is convex, *Linear and Multilinear Algebra* **56**(2008), 65–67.

CHI-KWONG LI, DEPARTMENT OF MATHEMATICS, COLLEGE OF WILLIAM AND MARY, WILLIAMSBURG, VIRGINIA 23185, U.S.A.

E-mail address: ckli@math.wm.edu

YIU-TUNG POON, DEPARTMENT OF MATHEMATICS, IOWA STATE UNIVERSITY, AMES, IOWA 50011, U.S.A.

E-mail address: ytpoon@iastate.edu

Received December 20, 2008.