

A BIRKHOFF TYPE TRANSITIVITY THEOREM FOR NON-SEPARABLE COMPLETELY METRIZABLE SPACES WITH APPLICATIONS TO LINEAR DYNAMICS

ANTONIOS MANOUSSOS

Communicated by Nikolai K. Nikolski

ABSTRACT. In this note we prove a Birkhoff type transitivity theorem for continuous maps acting on non-separable completely metrizable spaces and we give some applications for dynamics of bounded linear operators acting on complex Fréchet spaces. Among them we show that any positive power and any unimodular multiple of a topologically transitive linear operator is topologically transitive, generalizing similar results of S.I. Ansari and F. León-Saavedra–V. Müller for hypercyclic operators.

KEYWORDS: *Topological transitivity, hypercyclicity, Birkhoff's transitivity theorem, J -set, almost topological transitivity.*

MSC (2010): Primary 47A16, 54H20; Secondary 37B99, 54H15.

1. INTRODUCTION AND BASIC NOTIONS

The scope of this paper is to provide a tool for studying topologically transitive operators acting on non-separable Fréchet spaces by using techniques and known results from the theory of hypercyclic operators. This tool is the following theorem in which we show that each vector in the underlying space X is contained in "many" closed invariant subspaces such that the restriction of the operator to each one of them is hypercyclic:

THEOREM (Theorem 3.1). *Let $T : X \rightarrow X$ be a topologically transitive operator acting on a completely metrizable vector space X and let y be a vector of X . There exists a dense G_δ subset D of X such that for each $z \in D$ there exists a T -invariant (separable) closed subspace Y_z of X with $y, z \in Y_z$ such that the restriction of T to Y_z , $T : Y_z \rightarrow Y_z$, is hypercyclic.*

The previous theorem is derived from a Birkhoff type transitivity theorem for topologically transitive continuous selfmaps of a (not necessarily separable) completely metrizable space. More precisely we show the following:

THEOREM (Theorem 2.4). *Let X be a completely metrizable space which has no isolated points. Let $T : X \rightarrow X$ be a continuous map acting on X and let $x \in X$. If T is topologically transitive there exists a dense G_δ subset D of X with the following properties:*

(i) *Every point $z \in D$ is recurrent, that is there exists a strictly increasing sequence of positive integers $\{k_n\}$ such that $T^{k_n}z \rightarrow z$.*

(ii) *The point x belongs to the orbit closure $\overline{O(z, T)}$ of z for every $z \in D$; in particular x belongs to the limit set $L(z)$ of z for every $z \in D$, i.e. for each $z \in D$ there exists a strictly increasing sequence of positive integers $\{m_n\}$ such that $T^{m_n}z \rightarrow x$.*

This result comes by "localizing" Birkhoff's transitivity theorem; see Theorem 2.2 and Corollary 2.3. This local behavior was hidden in the proof of Birkhoff's transitivity theorem behind the use of a countable base of X . Recall that a continuous map $T : X \rightarrow X$ acting on a Hausdorff topological space X is called *topologically transitive* if for every pair of non-empty open subsets U, V of X there exists a non-negative integer n such that $T^n U \cap V \neq \emptyset$. In case X is completely metrizable and separable, Birkhoff's transitivity theorem [7] states that topological transitivity is equivalent to *hypercyclicity*, i.e. there exists a point x in X which has a dense orbit. The *orbit* of $x \in X$ under T is the set $O(x, T) := \{T^n x : n \in \mathbb{N} \cup \{0\}\}$. Hypercyclic operators on separable Fréchet spaces have been a subject of extensive study in the last decades. Very good references are the recent books [4] and [11].

The applications we give in Section 3 indicate the possibility to use the previous mentioned results to work on non-separable Fréchet spaces by applying technics and known results from the theory of hypercyclic operators. Namely, in Section 3, we show the following theorem which generalizes similar results of S.I. Ansari [1] and F. León-Saavedra-V. Müller [12] for the case of a hypercyclic operator:

THEOREM (Theorem 3.2). *Let $T : X \rightarrow X$ be a topologically transitive operator acting on a complex Fréchet space X and let $x \in X$. Then*

(i) *The operator $T^p : X \rightarrow X$ is topologically transitive for every positive integer p and there exists a dense subset D of X with the following properties:*

(a) *Every vector $z \in D$ is a recurrent vector for the operator T^p .*

(b) *The vector x belongs to the limit set $L_{T^p}(z)$ for every positive integer p and for every $z \in D$.*

(ii) *If λ is a complex number of modulus 1 the operator $\lambda T : X \rightarrow X$ is topologically transitive and there exists a dense subset D of X with the following properties:*

(a) *Every vector $z \in D$ is a recurrent vector for the operator λT .*

(b) *The vector x belongs to the limit set $L_{\lambda T}(z)$ for every $|\lambda| = 1$ and for every $z \in D$.*

We finish Section 3 with a characterization, similar to that of H.N. Salas in [14], of topological transitivity of a backward unilateral weighted shift on $l_2(H)$,

where H is a (not necessarily separable) Hilbert space, in terms of its weight sequence; see Proposition 3.4.

Finally, in Section 4 we show that every continuous almost topologically transitive map acting on a completely metrizable space which has no isolated points is topologically transitive; see Proposition 4.3. Recall that a continuous map T acting on a Hausdorff topological space X is called *almost topologically transitive* if for every pair of non-empty open sets $U, V \subset X$ there exists a non-negative integer n such that $T^n U \cap V \neq \emptyset$ or $T^n V \cap U \neq \emptyset$.

2. A BIRKHOFF TYPE TRANSITIVITY THEOREM FOR NON-SEPARABLE COMPLETELY METRIZABLE SPACES

Birkhoff's transitivity theorem can be reformulated using some very useful concepts from the theories of dynamical systems and topological transformation groups, namely the concept of the *limit set* of a point $x \in X$:

$$L(x) = \{y \in X : \text{there exists a strictly increasing sequence} \\ \text{of positive integers } \{k_n\} \text{ such that } T^{k_n} x \rightarrow y\}$$

that describes the limit behavior of the orbit $O(x, T)$ and the generalized (prolongational) limit set of $x \in X$ that describes the asymptotic behavior of the orbits of nearby points to $x \in X$:

$$J(x) = \{y \in X : \text{there exist a strictly increasing sequence of positive} \\ \text{integers } \{k_n\} \text{ and a sequence } \{x_n\} \subset X \text{ such that } x_n \rightarrow x \text{ and } T^{k_n} x_n \rightarrow y\}.$$

The limit and the extended limit sets have their roots in the stability theory of dynamical systems in which they are mainly used to describe the Lyapunov and the asymptotic stability of an equilibrium point. They are T -invariant and closed subsets of X , see e.g. Proposition 2.6 of [9]. In view of the concept of limit sets, a bounded linear operator $T : X \rightarrow X$ acting on a separable Fréchet space X is hypercyclic if and only if $L(x) = X$ for some non-zero vector $x \in X$. And Birkhoff's transitivity theorem says that T is hypercyclic if and only if $J(x) = X$ for every $x \in X$.

REMARK 2.1. If X is a completely metrizable space which has no isolated points (hence X is uncountable), the orbit of a point $x \in X$ (which is a countable set) is dense in X if and only if $L(x) = X$. In a similar way, topological transitivity, i.e. $J(x) \cup O(x, T) = X$ for every $x \in X$, means that $J(x) = X$ for every $x \in X$ since $J(x)$ is a closed subset of X . In general, if X is a completely metrizable space, one can use instead of the limit set $L(x)$ the orbit closure $\overline{O(x, T)} = O(x, T) \cup L(x)$ and instead of the generalized limit set $J(x)$ the union $O(x, T) \cup J(x) =: D(x)$ (the concept of the D -set comes also from the stability theory of dynamical systems and it is called the prolongation of the orbit $O(x, T)$, see e.g. [6]).

The following theorem, which can be seen as a "localized" Birkhoff's transitivity theorem, plays a key role in our approach to topological transitivity. The symbol $B(x, \varepsilon)$ stands for the open ball centered at x with radius $\varepsilon > 0$. Recall that a point $x \in X$ is called *recurrent* if $x \in L(x)$.

THEOREM 2.2. *Let X be a completely metrizable space and let $T : X \rightarrow X$ be a continuous map acting on X . Assume that there are points $x, y \in X$ and a positive number $\varepsilon > 0$ such that $x \in J(w)$ for every $w \in B(y, \varepsilon)$. Then*

- (i) *there is a point $z \in B(y, \varepsilon)$ such that $x \in L(z)$;*
- (ii) *if, moreover, every point $w \in B(y, \varepsilon)$ is non-wandering, i.e. $w \in J(w)$, there is a recurrent point $z \in B(y, \varepsilon)$ such that $x \in L(z)$.*

Proof. We give only the proof of claim (ii) since the proof of claim (i) is similar. Since $x \in J(y)$ there exist a point $y_1 \in B(y, \varepsilon)$, a positive integer k_1 and an open ball $B(y_1, \varepsilon_1) \subset B(y, \varepsilon)$, with $\varepsilon_1 < 1$ such that $T^{k_1}B(y_1, \varepsilon_1) \subset B(x, 1)$. And since $y_1 \in J(y_1)$ there exist a point $w_1 \in B(y_1, \varepsilon_1)$, an open ball $B(w_1, r_1) \subset B(y_1, \varepsilon_1)$, for some $r_1 > 0$, and a positive integer m_1 such that $T^{m_1}B(w_1, r_1) \subset B(y_1, \varepsilon_1)$. Note that $x \in J(w_1)$ since $x \in J(w)$ for every $w \in B(y, \varepsilon)$. Thus, there exist a point $y_2 \in B(w_1, r_1)$, a positive integer $k_2 > k_1$ and an open ball $B(y_2, \varepsilon_2) \subset B(w_1, r_1)$ with $\varepsilon_2 < 1/2$ such that $T^{k_2}B(y_2, \varepsilon_2) \subset B(x, 1/2)$. Now we can proceed as before. Since $y_2 \in J(y_2)$ there exist a point $w_2 \in B(y_2, \varepsilon_2)$, an open ball $B(w_2, r_2) \subset B(y_2, \varepsilon_2)$, for some $r_2 > 0$, and a positive integer $m_2 > m_1$ such that $T^{m_2}B(w_2, r_2) \subset B(y_2, \varepsilon_2)$. Proceeding by induction we can find two sequences of points $\{y_n\}$ and $\{w_n\}$, two strictly increasing sequences of positive integers $\{k_n\}$ and $\{m_n\}$ and two sequences of positive numbers $r_n \leq \varepsilon_n < 1/2^n$ with the following properties:

- (a) $B(y_{n+1}, \varepsilon_{n+1}) \subset B(w_n, r_n) \subset B(y_n, \varepsilon_n)$,
- (b) $T^{k_{n+1}}B(y_n, \varepsilon_n) \subset B(x, 1/(n+1))$ and
- (c) $T^{m_n}B(w_n, r_n) \subset B(y_n, \varepsilon_n)$,

for every $n \in \mathbb{N}$. Since X is a complete metric space and $r_n \leq \varepsilon_n < 1/2^n$ then

$$\bigcap_n B(y_n, \varepsilon_n) = \bigcap_n B(w_n, r_n) = \{z\}, \quad \text{for some } z \in X.$$

Therefore, $T^{k_n}z \rightarrow x$ by property (b) and $T^{m_n}z \rightarrow z$ by properties (a) and (c). ■

COROLLARY 2.3. *Let (X, d) be a complete metric space with metric d and let $T : X \rightarrow X$ be a continuous map acting on X . Assume that there are points $x, y \in X$ and a positive number $\varepsilon > 0$ such that $x \in J(w)$ for every $w \in B(y, \varepsilon)$. Then*

- (i) *There is a dense G_δ subset D of $\overline{B(y, \varepsilon)}$ such that $x \in L(z)$ for every $z \in D$.*
- (ii) *If, moreover, every point in $B(y, \varepsilon)$ is non-wandering, i.e. $w \in J(w)$, there is a dense G_δ subset D of $\overline{B(y, \varepsilon)}$ such that $x \in L(z)$ and z is recurrent, for every $z \in D$.*

Proof. By a diagonal procedure, see Lemma 2.4 of [8], $x \in J(w)$ for every $w \in \overline{B(y, \varepsilon)}$ since $x \in J(w)$ for every $w \in B(y, \varepsilon)$. Moreover, every point of

the closure of $B(y, \varepsilon)$ is non-wandering. We do that in order to apply the Baire category theorem and Theorem 2.2 for the complete metric space $\overline{B(y, \varepsilon)}$. Now, in (i), let $D := \{z \in \overline{B(y, \varepsilon)}$ such that $x \in L(z)\}$. Then D can be written as a countable intersection of open and dense subset of $\overline{B(y, \varepsilon)}$ in the following way:

$$D = \bigcap_{s, n \in \mathbb{N}} \bigcup_{m > n} T^{-m}B(x, \frac{1}{s}) \cap \overline{B(y, \varepsilon)}.$$

Similarly, in (ii) let $D := \{z \in \overline{B(y, \varepsilon)}$ such that $x \in L(z)$ and $z \in L(z)\}$. Then

$$D = \bigcap_{s, n \in \mathbb{N}} \left(\bigcup_{m > n} T^{-m}B(x, \frac{1}{s}) \cap \bigcup_{l > n} \left\{ w \in X : d(T^l w, w) < \frac{1}{s} \right\} \right) \cap \overline{B(y, \varepsilon)}.$$

Hence, in both cases (i) and (ii), the set D is a dense G_δ subset of $\overline{B(y, \varepsilon)}$. ■

The following two results are immediate corollaries of Theorem 2.2. They can also be found in Satz 1.2.2 of [10] and in a more general form in Theorem 26 of [3]. We are grateful to K.-G. Grosse-Erdmann for pointing out this to us.

THEOREM 2.4 (A Birkhoff type transitivity theorem for non-separable completely metrizable spaces). *Let X be a completely metrizable space which has no isolated points. Let $T : X \rightarrow X$ be a continuous map acting on X and let $x \in X$. If T is topologically transitive there exists a dense G_δ subset D of X with the following properties:*

- (i) *Every point $z \in D$ is recurrent.*
- (ii) *The point x belongs to the limit set $L(z)$ for every $z \in D$.*

COROLLARY 2.5. *Let X be a completely metrizable space which has no isolated points. Let $T : X \rightarrow X$ be a continuous map acting on X and let A be a countable subset of X . If T is topologically transitive there exists a dense G_δ subset D of X with the following properties:*

- (i) *Every point $z \in D$ is recurrent.*
- (ii) *The closure \overline{A} of A is a subset of $L(z)$ for every $z \in D$.*

Note that if X is completely metrizable and separable then we can recover the original transitivity theorem of Birkhoff by applying the previous corollary.

3. APPLICATIONS TO LINEAR DYNAMICS

In this section we will use Theorem 3.1 and known results from the theory of hypercyclic operators to derive similar results for the non-separable case.

We quote from the Preface of [11], "... Some of the deepest, most beautiful and most useful results from linear dynamics is Ansari's theorem on the powers of hypercyclic operators and the León–Müller theorem on the hypercyclicity of unimodular multiples of hypercyclic operators ...". More precisely, Ansari showed in [1] that if T is a hypercyclic operator then any positive power of T is

also a hypercyclic operator with the same set of hypercyclic vectors. And León-Saavedra and Müller showed in [12] that if T is a hypercyclic operator and λ is a complex number of modulus 1 then λT is also a hypercyclic operator with the same hypercyclic vectors. The reason that we can obtain similar results in the more general setting of a topologically transitive operator is because each vector in X is contained in "many" closed invariant subspaces such that the restriction of the operator to each one of them is hypercyclic:

THEOREM 3.1. *Let $T : X \rightarrow X$ be a topologically transitive operator acting on a completely metrizable vector space X and let y be a vector of X . There exists a dense G_δ subset D of X such that for each $z \in D$ there exists a T -invariant (separable) closed subspace Y_z of X with $y, z \in Y_z$ such that the restriction of T to Y_z , $T : Y_z \rightarrow Y_z$, is hypercyclic.*

Proof. Let W_0 denote the closed linear span of the orbit $O(y, T)$, that is W_0 is the closure of the subspace $\{P(T)y : P \text{ polynomial}\}$. Since T is topologically transitive then, by Corollary 2.5, there exists a dense G_δ subset D of X which consists of recurrent vectors such that $W_0 \subset \overline{O(x, T)}$ for each $x \in D$. Now, choose a vector $z \in D$ and set $x_1 := z$. Thus, $W_0 \subset \overline{O(x_1, T)}$ and $x_1 \in L(x_1)$. Let W_1 be the closed linear span of the orbit $O(x_1, T)$. By the same corollary there exists a vector $x_2 \in X$ such that $W_0 \subset \overline{O(x_1, T)} \subset W_1 \subset \overline{O(x_2, T)}$ and $x_2 \in L(x_2)$. Proceeding by induction, there is a sequence of vectors $\{x_n\}_{n \in \mathbb{N}}$ in X such that $W_{n-1} \subset \overline{O(x_n, T)} \subset W_n$, where W_n denotes the closed linear span of the orbit $O(x_n, T)$, and $x_n \in L(x_n)$ for every $n \in \mathbb{N}$. Set $Y_z := \bigcup_{n=0}^{+\infty} W_n$. Obviously, Y_z is a T -invariant, closed and separable subspace of X . Now, let us show that the restriction of T to Y_z , $T : Y_z \rightarrow Y_z$, is hypercyclic. Let $U, V \subset Y_z$ be two non-empty relatively open subsets of Y_z . Since $Y_z := \bigcup_{n=1}^{+\infty} \overline{O(x_n, T)}$ and $\overline{O(x_n, T)} \subset \overline{O(x_{n+1}, T)}$ for every $n \in \mathbb{N}$, there exist a positive integer p and some non-negative integers k and m such that $T^k x_p \in U$ and $T^m x_p \in V$ (note that the vector x_p is common for both U and V). Since x_p is a recurrent vector and V is open in T_z we may assume that $m > k$. Now note that $T^{m-k}(T^k x_p) = T^m x_p \in V$. Thus, $T^{m-k}U \cap V \neq \emptyset$ and hence $T : Y_z \rightarrow Y_z$ is topologically transitive. Therefore, by Birkhoff's transitivity theorem, $T : Y_z \rightarrow Y_z$ is hypercyclic and since z was an arbitrary vector of D , which is a dense G_δ subset of X , the proof is completed. ■

Now we are ready to show Ansari's and León-Müller theorems for topologically transitive operators:

THEOREM 3.2. *Let $T : X \rightarrow X$ be a topologically transitive operator acting on a complex Fréchet space X and let $x \in X$. Then*

(i) *The operator $T^p : X \rightarrow X$ is topologically transitive for every positive integer p and there exists a dense subset D of X with the following properties:*

(a) *Every vector $z \in D$ is a recurrent vector for the operator T^p .*

(b) *The vector x belongs to the limit set $L_{T^p}(z)$ for every positive integer p and*

for every $z \in D$.

(ii) If λ is a complex number of modulus 1 the operator $\lambda T : X \rightarrow X$ is topologically transitive and there exists a dense subset D of X with the following properties:

(a) Every vector $z \in D$ is a recurrent vector for the operator λT .

(b) The vector x belongs to the limit set $L_{\lambda T}(z)$ for every $|\lambda| = 1$ and for every $z \in D$.

Proof. Let U, V be two non-empty open subsets of X and let $x \in U$. By Theorem 3.1 there is a vector $z \in V$ and a T -invariant separable closed subspace Y_z of X with $x, z \in Y_z$ such that the restriction of T to Y_z , $T : Y_z \rightarrow Y_z$, is hypercyclic. Hence, we can apply the original theorems of Ansari and León-Müller for the restriction $T : Y_z \rightarrow Y_z$. Thus, the operators $T^p : Y_z \rightarrow Y_z$ and $\lambda T : Y_z \rightarrow Y_z$ are hypercyclic and share the same set of hypercyclic vectors with $T : Y_z \rightarrow Y_z$. So we can find a (common) hypercyclic vector $w \in Y_z$ for the operators $T^p : Y_z \rightarrow Y_z$ and $\lambda T : Y_z \rightarrow Y_z$ such that the vectors x, z belong to the limit sets $L_{T^p}(w)$ and $L_{\lambda T}(w)$ respectively. Therefore, like in the proof of Theorem 3.1, the operators $T^p : X \rightarrow X$ and $\lambda T : X \rightarrow X$ are topologically transitive. In both cases (i) and (ii), let D be the union of the sets of hypercyclic vectors of the family of operators $T : Y_z \rightarrow Y_z$ for z in a dense G_δ subset of X . Thus, D is dense in X . ■

REMARK 3.3. Linearity of the map under consideration is essential in the proof of Theorem 3.2. It is evident, that even for real piecewise linear maps, a power of a topologically transitive map may not be topologically transitive; see e.g. [2].

We finish this section with a characterization, similar to that of H.N. Salas in [14], of topological transitivity of a backward unilateral weighted shift on $l_2(H)$, where H is a (not necessarily separable) Hilbert space, in terms of its weight sequence. The motivation of this problem comes from a work of T. Bermúdez and N.J. Kalton [5]. In this paper they showed that spaces like $l^\infty(\mathbb{N})$ and $l^\infty(\mathbb{Z})$ do not support topologically transitive operators. On the other hand, following an example suggested by J.H. Shapiro, they showed that every non-separable Hilbert space supports a topologically transitive operator. To do that they wrote H as $l_2(X)$ for some Hilbert space X of the same density character with H , i.e. a space with the same least cardinality of a dense subset. Then they showed that the operator $2B$, where B is the backward unilateral shift on $l_2(X)$, is topologically transitive. Hence, a problem that naturally arises is to give a characterization of topological transitivity of a backward unilateral weighted shift on $l_2(H)$, where H is a (not necessarily separable) Hilbert space, in terms of its weight sequence.

PROPOSITION 3.4. Let H be Hilbert space and let $T : l_2(H) \rightarrow l_2(H)$ be a unilateral backward weighted shift with positive weight sequence (w_n) . The following are equivalent:

(i) T is topologically transitive.

(ii) *There exists a non-trivial T -invariant (separable) closed subspace $Y \subset l_2(H)$ on which the restriction of T to Y , $T : Y \rightarrow Y$, is hypercyclic.*

(iii) *The restriction $T : Y \rightarrow Y$ to any T -invariant (separable) closed subspace $Y \subset l_2(H)$ is hypercyclic.*

(iv) $\limsup_{n \rightarrow \infty} (w_1 \cdots w_n) \rightarrow +\infty$.

Proof. Property (i) implies (ii) by Theorem 3.1. From Salas characterization of hypercyclic unilateral backward shifts in [14], claim (ii) implies (iii). From the same theorem property (iv) implies (iii) and vice versa. Finally, it is easy to verify that claim (iii) implies (i). ■

4. ALMOST TOPOLOGICALLY TRANSITIVE OPERATORS ARE TOPOLOGICALLY TRANSITIVE

Among the concepts that generalize, or are closely related, to topological transitivity is the concept of a topological semi-transitive operator; see e.g. [13] for the case of operator algebras. Recall that an operator $T : X \rightarrow X$ acting on a topological vector space X is called *topologically semi-transitive* if for every two non-zero vectors $x, y \in X$ we have that $x \in \overline{O(y, T)}$ or $y \in \overline{O(x, T)}$. There are some disadvantages in this concept for the case of the action of a semigroup of linear operators. One disadvantage is that topological semi-transitivity is not a topological concept. Another disadvantage is that topological semi-transitivity is very restrictive, since even a chaotic operator is not topologically semi-transitive (to see that just consider two non-zero periodic vectors). A concept that looks more attractive to generalize topological transitivity is the concept of almost topological transitivity:

DEFINITION 4.1. Let X be a Hausdorff topological space and let $T : X \rightarrow X$ be a continuous map acting on X . We call T *almost topologically transitive* if for every pair of non-empty open sets $U, V \subset X$ there exists a non-negative integer n such that $T^n U \cap V \neq \emptyset$ or $T^n V \cap U \neq \emptyset$. That is $x \in D(y) := O(y, T) \cup J(y)$ or $y \in D(x) := O(x, T) \cup J(x)$ for every $x, y \in X$.

Note that if X is a completely metrizable space which has no isolated points the orbit $O(x, T)$ has empty interior for every $x \in X$. In this case the following proposition shows that a continuous map $T : X \rightarrow X$ is almost topologically transitive if $x \in J(y)$ or $y \in J(x)$ for every $x, y \in X$.

PROPOSITION 4.2. *Let X be a completely metrizable space which has no isolated points. A continuous map $T : X \rightarrow X$ is almost topologically transitive if $x \in J(y)$ or $y \in J(x)$ for every $x, y \in X$.*

Proof. We proceed by contradiction. Assume that there exists a pair of (not necessarily distinct) vectors $x, y \in X$ such that $x \notin J(y)$ and $y \notin J(x)$. Hence, $x \in O(y, T)$ or $y \in O(x, T)$ since T is almost topologically transitive. Without loss

of generality we may assume that $x \in O(y, T)$. Since the orbit $O(y, T)$ has empty interior there is a sequence $\{x_n\}_{n \in \mathbb{N}}$ in X such that $x_n \rightarrow x$ and $x_n \notin O(y, T)$ for every $n \in \mathbb{N}$. Thus, the following hold: (a) $x_n \in J(y)$ for infinitely many $n \in \mathbb{N}$, so $x \in J(y)$ since $J(y)$ is a closed subset of X which is a contradiction or (b) $y \in D(x_n) := O(x_n, T) \cup J(x_n)$ eventually. In that case, using a diagonal procedure similar to the one in Lemma 2.4 of [8], it follows that $y \in D(x) := O(x, T) \cup J(x)$. This implies that $y \in O(x, T)$ since $y \notin J(x)$. But also $x \in O(y, T)$, hence $O(x, T) = O(y, T)$. So, if x, y are distinct vectors then $O(x, T) = O(y, T)$ is a periodic orbit, thus $x \in J(y)$ (and $y \in J(x)$) which is a contradiction. If $x = y$, and since $x_n \notin O(y, T) = O(x, T)$, we may assume that $x_n \neq x$ for every $n \in \mathbb{N}$. Hence, as we showed above for the case of two distinct points, $x_n \in J(x)$ for infinitely many $n \in \mathbb{N}$ or $x \in J(x_n)$ eventually. In both cases, $x \in J(x) = J(y)$, by Lemma 2.4 of [8], which is again a contradiction. ■

The above proposition and Corollary 2.3 imply the following:

PROPOSITION 4.3. *Let X be a completely metrizable space which has no isolated points and let $T : X \rightarrow X$ be an almost topologically transitive selfmap of X . Then T is topologically transitive.*

Proof. Note that $x \in J(x)$ for every $x \in X$ since T is almost topologically transitive. Now, let us show that $J(x) = X$ for every $x \in X$. We argue by contradiction. Assume that $J(x) \neq X$ for some $x \in X$. Hence, $x \in J(y)$ for every y is in the complement of $J(x)$. Therefore, by Corollary 2.3, there exist a point $y \notin J(x)$ and two strictly increasing sequences of positive integers $\{k_n\}$ and $\{m_n\}$ such that $T^{k_n}y \rightarrow x$ and $T^{m_n}y \rightarrow y$. We may assume that $m_n - k_n \rightarrow +\infty$. Thus, $T^{m_n - k_n}(T^{k_n}y) = T^{m_n}y \rightarrow y$ and $T^{k_n}y \rightarrow x$, hence $y \in J(x)$ which is a contradiction. ■

Acknowledgements. We would like to thank Frédéric Bayart, George Costakis, Karl-Goswin Grosse-Erdmann and Alfredo Peris for their very useful comments.

During this research the author was fully supported by SFB 701 "Spektrale Strukturen und Topologische Methoden in der Mathematik" at the University of Bielefeld, Germany.

REFERENCES

- [1] S.I. ANSARI, Hypercyclic and cyclic vectors, *J. Funct. Anal.* **128**(1995), 374–383.
- [2] J. BANKS, Regular periodic decompositions for topologically transitive maps, *Ergodic Theory Dynam. Systems* **17**(1997), 505–529.
- [3] F. BAYART, K.-G. GROSSE-ERDMANN, V. NESTORIDIS, C. PAPADIMITROPOULOS, Abstract theory of universal series and applications, *Proc. London Math. Soc. (3)* **96**(2008), 417–463.

- [4] F. BAYART, É. MATHERON, *Dynamics of Linear Operators*, Cambridge Tracts in Math., vol. 179, Cambridge Univ. Press, Cambridge 2009.
- [5] T. BERMÚDEZ, N.J. KALTON, The range of operators on von Neumann algebras, *Proc. Amer. Math. Soc.* **130**(2002), 1447–1455.
- [6] N.P. BHATIA, G.P. SZEGÖ, *Stability Theory of Dynamical Systems*, Grundlehren Math. Wiss., vol. 161, Springer-Verlag, New York-Berlin 1970.
- [7] G.D. BIRKHOFF, Surface transformations and their dynamical applications, *Acta Math.* **43**(1922), 1–119.
- [8] G. COSTAKIS, A. MANOUSSOS, J -class weighted shifts on the space of bounded sequences of complex numbers, *Integral Equations Operator Theory* **62**(2008), 149–158.
- [9] G. COSTAKIS, A. MANOUSSOS, J -class operators and hypercyclicity, *J. Operator Theory* **67**(2012), 101–119.
- [10] K.-G. GROSSE-ERDMANN, Holomorphe Monster und universelle Funktionen, *Mitt. Math. Sem. Giessen* **176**(1987).
- [11] K.-G. GROSSE-ERDMANN, A. PERIS, *Linear Chaos*, Universitext, Springer, London 2011.
- [12] F. LEÓN-SAAVEDRA, V. MÜLLER, Rotations of hypercyclic and supercyclic operators, *Integral Equations Operator Theory* **50**(2004), 385–391.
- [13] P. ROSENTHAL, V.G. TROITSKY, Strictly semi-transitive operator algebras, *J. Operator Theory* **53**(2005), 315–329.
- [14] H.N. SALAS, Hypercyclic weighted shifts, *Trans. Amer. Math. Soc.* **347**(1995), 993–1004.

ANTONIOS MANOUSSOS, FAKULTÄT FÜR MATHEMATIK, SFB 701, UNIVERSITÄT BIELEFELD, POSTFACH 100131, D-33501 BIELEFELD, GERMANY
E-mail address: amanouss@math.uni-bielefeld.de
URL: <http://www.math.uni-bielefeld.de/~amanouss>

Received May 12, 2011; revised January 20, 2013.